

Background independent quantizations: the scalar field II

Wojciech Kamiński^{1,*}, Jerzy Lewandowski^{1,2,†}, Andrzej Okołów^{1,3,‡}

1. *Instytut Fizyki Teoretycznej, Uniwersytet Warszawski, ul. Hoża 69, 00-681 Warszawa, Poland*
2. *Perimeter Institute for Theoretical Physics, 31 Caroline Street North, Waterloo, Ontario N2L 2Y5, Canada*
3. *Department of Physics and Astronomy, Louisiana State University, Baton Rouge, LA 70803-4001*

Abstract

We are concerned with the issue of quantization of a scalar field in a diffeomorphism invariant manner. We apply the method used in Loop Quantum Gravity. It relies on the specific choice of scalar field variables referred to as the polymer variables. The quantization, in our formulation, amounts to introducing the ‘quantum’ *polymer $\ast$ -star algebra* and looking for positive linear functionals, called states. Assumed in our paper homeomorphism invariance allows to derive the complete class of the states. They are determined by the homeomorphism invariant states defined on the CW-complex $\ast$ -algebra. The corresponding GNS representations of the polymer $\ast$ -algebra and their self-adjoint extensions are derived, the equivalence classes are found and invariant subspaces characterized. In the preceding letter (the part I) we outlined those results. Here, we present the technical details.

1 Introduction

1.1 Motivation

Einstein’s theory of gravity coupled (or not) to a matter field is a prominent example of a so called “background independent” theory. The phrase “background independent” means that the theory is defined on a bare manifold endowed with no geometry or affine structure. In this case, it is natural to look for a corresponding quantum theory that is

^{*}wkaminsk@fuw.edu.pl

[†]lewand@fuw.edu.pl

[‡]oko@fuw.edu.pl

also manifestly background independent (see [1] for a profound discussion of that issue). It is conceivable, that a background independent quantum theory can even be derived from a background *dependent* framework. Another possibility is, that the classical limit of a given quantum theory (the classical GR for example) has more symmetries (the diffeomorphisms) than the underlying quantum theory. Therefore, we are not in the position to claim that every background dependent approach to a quantization of a background independent classical theory is wrong. Nonetheless, the first thing one should do is to try without introducing extra structures. Loop Quantum Gravity is the only known example. That name comes from the idea [2, 3, 4] of using field variables labeled by loops, typically non-intersecting but knotted. Later, embedded *graphs* were found a correct tool [5, 6, 7] which gave rise to the present form of the theory. Therefore, there were attempts to promote the theory under a new name like “Quantum Geometry” [8]-[12] or “Quantum Spin Dynamics” [16]-[18]. LQG [19] is a canonical theory, it relies on the $3+1$ decomposition of space-time into the ‘space’ M and ‘time’ $\mathbb{R}$. It is invariant with respect to the diffeomorphisms (piecewise analytic) of M . It concerns the space-time geometry and its dynamics, as well as coupled matter fields [20]. The matter fields however, have to be quantized in a new way, consistent with the LQG and background independent quantization. And this is the point we want to focus on in this work. We are concerned here with the background independent, canonical quantization of the scalar field. The first example of a background independent quantization of the scalar field was proposed by Thiemann in his pioneering work [20]. It was improved and analyzed by various authors [21]. In the current paper we systematically derive a broader class of examples. We show that our class is complete if certain simple assumptions, like the topological invariance, are made. In fact we consider only the GNS representations (and their self-adjoint extensions) defined by invariant states introduced on the appropriate $*$ -algebra. Among the constructed representations we identify: those which give essentially self adjoint operators, the equivalency classes, the irreducible representations. Each of them can be applied to the scalar field interacting with the quantum geometry in the frame of LQG. A brief discussion of our results is already published in the form of a letter [22].

Here is an outline of the work.

In Sec. 2 we introduce the polymer $*$ -algebra $\mathfrak{A}$, the key object studied in this work. It is constructed by the canonical quantization of a classical field ϕ and the canonically conjugate momentum π , both living on a manifold M . The first step is choosing basic field-variables. Briefly speaking (see Sec. 2.2 for the details) they are: the ‘position variables’ $e^{ik\phi(x)}$ (denoted $h_{k,x}$) and the ‘momentum variables’ $\int \pi f$ (denoted $\pi(f)$), where $k \in \mathbb{R}$ and f is a smearing function. The vector space of the smearing functions $\mathcal{F}$ is fixed in this section arbitrarily. Its specific choice will play a crucial role later. The basic variables together with the Poisson bracket form a Lie algebra, still classical. The polymer $*$ -algebra is defined (Sec. 2.3) by ‘putting hats’, meaning it is the quantum

enveloping algebra of the Lie algebra in question. It still depends on the choice of the smearing function space $\mathcal{F}$. Next, we recall the definition of a state on a $*$ -algebra and elements of the GNS construction. The conditions that will be imposed on the states are formulated in In Secs. 2.5 and 2.6. Briefly, they are: (i) the diffeomorphism (Sec. 3) and the homeomorphism (Sec. 4-9.1) invariance, and (ii) an extra condition ensuring that for every point x , the quantum operators $\rho(\hat{h}_{k,x})$, $k \in \mathbb{R}$, form a 1-dimensional group of unitary operators.

The first example of the polymer $*$ -algebra considered in this work is given by choosing for the smearing functions all the $C^{(n)}$, $n = 0, 1, \dots, \dots\infty$, compactly supported functions on the manifold M (Sec. 3). In this case we show there is exactly one $C^{(n)}$ -diffeomorphisms (homeomorphism in the $n = 0$ case) invariant state on the polymer $*$ -algebra $\mathfrak{A}$. The proof is a simplified version of the proof used in [23] in the case of the holonomy-flux $*$ -algebra. This state coincides with the one used thus far in LQG [20, 21].

The proof of that first result is very sensitive on the differentiability class of the smearing functions being equal or greater then the differentiability of the diffeomorphisms. The question that arises naturally, is: how much the uniqueness proved in Sec. 3 depends on those differentiability assumptions? Another natural choice for the smearing functions -also used in LQG- is the characteristic functions of bounded regions in M . The study of this case is the subject of the main part of this work.

Specifically, the space $\mathcal{F}_S$ of the smearing functions we consider throughout Secs. 4 - 9 is spanned by the characteristic functions of ball-like regions in the manifold M . The manifold is endowed with a piecewise-analytic structure and the regions are assumed to be piecewise-analytic simplexes (see Appendix). First (Sec. 4), we study the abelian algebra $\exp(\odot\mathcal{F}_S)$ freely generated by the characteristic functions, and its complexification called here the CW-complex $*$ -algebra. That algebra, well defined on its own, can be identified with the subalgebra of the $*$ -polymer algebra generated by all the momentum variables. We find all the homeomorphism invariant states on the CW-complex $*$ -algebra. They can be labeled by the states $\mu : \mathbb{C}[\tau] \rightarrow \mathbb{C}$ where $\mathbb{C}[\tau]$ is the $*$ -algebra of the polynomials of one real variable.

Every homeomorphism invariant state on the CW-complex $*$ -algebra admits a natural extension to the polymer $*$ -algebra $\mathfrak{A}$ (Sec. 5). The resulting state defined on $\mathfrak{A}$ is also homeomorphism invariant. In this way a 1-1 correspondence between the states $\mu : \mathbb{C}[\tau] \rightarrow \mathbb{C}$ and all the homeomorphism invariant states on $\mathfrak{A}$ which satisfy Property 2.10 (mentioned above) was established.

In the remaining part of the work (Secs. 6 - 9) we are concerned with the properties of the GNS representations corresponding to the homeomorphism invariant states on $\mathfrak{A}$ derived in Sec. 4 and their and self-adjoint extensions. An explicit form of the GNS representation defined by a state found in Sec. 4 is derived in Sec. 6. The issue of the self-adjointness of the resulting operators is studied in Sec 7. In particular,

we characterize all those representations which admit a unique self-adjoint extension according to Schmudgen [24]. The equivalence problem for the self-adjoint extensions of our representations is solved in Sec. 8. The reducibility issue is solved in Sec. 9. The results are summarized in Sec. 10.

The problem we studied is also relevant for the issue of the uniqueness of the diffeomorphism invariant state of quantum geometry in LQG. We elaborate on that in Sec. 10.

2 The polymer *-algebra $\mathfrak{A}$ and GNS construction

2.1 The classical scalar field

The classical scalar field in the canonical approach consists of a pair of fields (ϕ, π) defined on a N -real dimensional manifold M , where: $\phi \in C^{m_0}(M, \mathbb{R})$ where $C^{m_0}(M, \mathbb{R})$ stands for the space of the C^{m_0} real valued functions defined on M , $m_0 \in \mathbb{N}$ or $m_0 = \infty$, whereas π , called the canonical momentum, is a scalar density of the weight 1. The momentum π can be expressed by a function $\tilde{\pi} : U \rightarrow \mathbb{R}$ defined on an arbitrary region $U \subset M$ equipped with coordinates $(x^1, \dots, x^N)$. The function, however, depends on the coordinates in the following way: if $(x'^1, \dots, x'^N)$ is another coordinate system defined on U , then the corresponding momentum function $\tilde{\pi}'$ is such that at every $x \in U$

$$\tilde{\pi}(x)d^N x = \tilde{\pi}'(x)d^N x'. \quad (2.1)$$

The fields ϕ and π are called canonically conjugate in the sense of the Poisson bracket, usually wrote as

$$\{\phi(x), \tilde{\pi}(y)\} = \delta_x(y). \quad (2.2)$$

That Poisson bracket will be encoded in the Lie bracket of the Lie algebra introduced in the next subsection.

2.2 The classical polymer Lie algebra

Definition 2.1. *A polymer position variable is a (n exponentiated evaluation) function $h_{k,x} : C^{m_0}(M, \mathbb{R}) \rightarrow \mathbb{C}$ assigned to a pair $(k, x) \in \mathbb{R} \times M$ such that $k \neq 0$, and defined as follows*

$$h_{k,x} : \phi \mapsto e^{ik\phi(x)}. \quad (2.3)$$

The set of the polymer position variables is closed with respect to the complex conjugation

$$\overline{h_{k_i, x_i}} = h_{-k_i, x_i}. \quad (2.4)$$

Definition 2.2. *The polymer position variable space $\text{Cyl}_1^{\mathbb{R}}$ is the set of all the real valued linear combinations of all the polymer position variables, that is the set of all the linear combinations h such that can be written in the form*

$$h = \sum_{i=1}^n (a_i h_{k_i, x_i} + \bar{a}_i h_{-k_i, x_i}) \quad (2.5)$$

where $n \in \mathbb{N}$, $x_1, \dots, x_n \in M$, $k_1, \dots, k_n \in \mathbb{R} \setminus \{0\}$, $a_1, \dots, a_n \in \mathbb{C}$ are arbitrary.

We could well be using only real valued functions $\text{Re } h_{k,x}$ and $\text{Im } h_{k,x}$ up to this point. The complexification will come with the quantization.

A momentum variable will be assigned to a function $f : M \rightarrow \mathbb{R}$. It can be thought of as the integral $\int_M \pi f$. However, we will identify it with the operator defined by the Poisson brackets $\{h_{k,x}, \int_M \pi f\}$:

Definition 2.3. *A momentum variable is a linear map $\pi(f) : \text{Cyl}_1^{\mathbb{R}} \rightarrow \text{Cyl}_1^{\mathbb{R}}$ and defined by the following action on the polymer position variables*

$$\pi(f) h_{k,x} = i t f(x) h_{t,x}. \quad (2.6)$$

This definition is our version of the Poisson bracket (2.2). We fix a vector subspace $\mathcal{F}$ of the space of all the real valued functions $\text{Map}(M, \mathbb{R})$ defined on M and refer to it the space of the smearing functions. We consider only the momenta defined by the smearing functions $f \in \mathcal{F}$.

Definition 2.4. *The momentum variable space $\Pi_{\mathcal{F}}$ defined by a given space $\mathcal{F} \subset \text{Map}(M, \mathbb{R})$ of the smearing functions is the real vector space spanned by the linear maps $\pi(f)$ (2.6) such that $f \in \mathcal{F}$. The vector space $\mathcal{F}$ itself is referred to as the smearing functions space, and its elements as smearing functions.*

Actually, the map

$$\mathcal{F} \ni f \mapsto \pi(f) \in \Pi_{\mathcal{F}}, \quad (2.7)$$

is an isomorphism of the vector spaces.

Finally, we define:

Definition 2.5. *The polymer Lie algebra $(\mathfrak{A}_{\text{cl}}, \{\cdot, \cdot\})$ corresponding to a momentum variable space $\Pi_{\mathcal{F}}$ is the direct sum vector space*

$$\mathfrak{A}_{\text{cl}} := \text{Cyl}_1^{\mathbb{R}} \oplus \Pi_{\mathcal{F}}, \quad (2.8)$$

equipped with the following Lie bracket $\{\cdot, \cdot\}$:

$$\{(h, \pi), (h', \pi')\} := (\pi' h - \pi h', 0). \quad (2.9)$$

The Lie bracket $\{\cdot, \cdot\}$ encodes the structure of the Poisson bracket (2.2) and this is the reason we denote it in this way (rather than the usual $[\cdot, \cdot]$).

Later in this paper, we will consider the following two examples of the smearing functions space $\mathcal{F}$ and the corresponding of the momentum variable spaces:

Example:

1.

$$\mathcal{F} = C_0^{m_0}(M, \mathbb{R})$$

where the second subscript 0 stands for ‘compactly supported’.

2. $\mathcal{F}$ spanned by the characteristic functions of suitably defined family of open subsets of M

Our main results will concern the second example. But until the end of this section we keep the smearing functions space $\mathcal{F}$ general.

2.3 The quantum polymer *-algebra $\mathfrak{A}$

In order to turn the polymer position and momentum variables (Definition 2.1 and 2.3) into quantum operators we will construct from them a *-algebra. Briefly speaking, the commutation relations of the operators will be defined as i times the Lie bracket of the corresponding elements of the classical Lie algebra. To do it in an exact way we formulate a definition analogous to that of the universal enveloping algebra.

Let $(\mathfrak{A}_{\text{cl}}, \{\cdot, \cdot\})$ be an arbitrary real Lie algebra.

First, consider a complex Lie algebra $(\mathfrak{A}_{\text{cl}}^{\mathbb{C}}, \{\cdot, \cdot\})$, the complexification of $(\mathfrak{A}_{\text{cl}}, \{\cdot, \cdot\})$. There is the natural complex conjugation $\bar{\cdot} : \mathfrak{A}_{\text{cl}}^{\mathbb{C}} \rightarrow \mathfrak{A}_{\text{cl}}^{\mathbb{C}}$.

Next, consider the huge space

$$\bigoplus_{n=0}^{\infty} (\mathfrak{A}_{\text{cl}}^{\mathbb{C}})^{\otimes n} \tag{2.10}$$

where

$$(\mathfrak{A}_{\text{cl}}^{\mathbb{C}})^0 := \mathbb{C}. \tag{2.11}$$

It has the natural complex associative algebra structure defined by the complex vector space structure and the operation $\otimes$. There is also naturally defined anti-isomorphism-involution $\cdot^*$ in it, such that

$$(a_1 \otimes \cdots \otimes a_n)^* = \overline{a_n} \otimes \cdots \otimes \overline{a_1}. \tag{2.12}$$

Next, introduce the double sided ideal J generated by the following subset

$$\left\{ a \otimes b - b \otimes a - i\{a, b\} : a, b \in \mathfrak{A}_{\text{cl}}^{\mathbb{C}} \right\}. \tag{2.13}$$

Ideal J is preserved by the involution $\cdot^*$, hence $\cdot^*$ passes to the quotient $\bigoplus_{n=0}^{\infty} (\mathfrak{A}_{\text{cl}}^{\mathbb{C}})^{\otimes n} / J$ (and is denoted by the same symbol $\cdot^*$).

Finally, define:

Definition 2.6. *The quantum enveloping algebra $(\mathfrak{A}, \cdot^*)$ of a real Lie algebra $(\mathfrak{A}_{\text{cl}}, \{\cdot, \cdot\})$ is the associative, unital, $\cdot^*$ -algebra*

$$\mathfrak{A} = \bigoplus_{n=0}^{\infty} (\mathfrak{A}_{\text{cl}}^{\mathbb{C}})^{\otimes n} / J. \quad (2.14)$$

From now on, until the end of the paper we will be considering the quantum enveloping algebra of the polymer Lie algebra:

Definition 2.7. *A polymer $\cdot^*$ -algebra $\mathfrak{A}$ is the quantum enveloping algebra of a polymer Lie algebra.*

Elements of $\mathfrak{A}$ can be heuristically thought of as quantum operators corresponding to polynomials in the polymer position and momentum variables. (They become operators indeed, only given a representation of $\mathfrak{A}$ - see the next subsection).

We will be using the following notation: an element $[a_1 \otimes \cdots \otimes a_n] \in \mathfrak{A}$ corresponding to $a_1 \otimes \cdots \otimes a_n$, where $a_1, \dots, a_n \in \mathfrak{A}_{\text{cl}}^{\mathbb{C}}$ will be denoted by $\widehat{a_1} \dots \widehat{a_n}$,

$$\widehat{a_1} \dots \widehat{a_n} := [a_1 \otimes \cdots \otimes a_n]. \quad (2.15)$$

In particular, we will be denoting

$$\hat{\pi}(f) = [\pi(f)] \in \mathfrak{A}, \quad \hat{h}_{k,x} = [h_{k,x}] \in \mathfrak{A}, \quad (2.16)$$

and will be calling them the quantum momentum, and, respectively, quantum position variable.

The basic commutators $[A, B] = AB - BA$ in the polymer $\cdot^*$ -algebra $\mathfrak{A}$ are

$$[\hat{h}_{k,x}, \hat{\pi}(f)] = -kf(x)\hat{h}_{k,x}, \quad (2.17)$$

$$[\hat{h}_{t,x}, \hat{h}_{t',x'}] = 0 = [\hat{\pi}(f), \hat{\pi}(f)] \quad (2.18)$$

for arbitrary $x, x' \in M$, $k, k' \in \mathbb{R} \setminus \{0\}$ and $\pi(f) \in \Pi_{\mathcal{F}}$.

Owing to the commutation relations, every element of the polymer $\cdot^*$ -algebra $\mathfrak{A}$ corresponding to a momentum variable space $\Pi_{\mathcal{F}}$ can be written as a complex linear combination of the elements of the form

$$\prod_{i=1}^n \hat{h}_{k_i, x_i} \prod_{j=1}^m \hat{\pi}(f_j), \quad (2.19)$$

where $x_i \in M$, $k_i \neq 0$, and $\pi(f_j) \in \Pi_{\mathcal{F}}$, and the cases either $\prod_{i=1}^n \hat{h}_{k_i, x_i}$, or $\prod_{j=1}^m \hat{\pi}(f_j)$ are included as corresponding to $m = 0$ and, respectively, $n = 0$.

This decomposition is not necessarily unique, though. To begin with the factors h_{t_i, x_i} commute among themselves as well as the factors $\pi(f_i)$. The second source of the non-uniqueness, is the identities satisfied by the momentum variables Definition 2.3

$$\pi(a_1 f_1 + \dots + a_k f_k) = a_1 \pi(f_1) + \dots + a_k \pi(f_k) \quad (2.20)$$

for every finite set of functions $f_1, \dots, f_k \in \mathcal{F}$ and real numbers $a_1, \dots, a_k$. On the other hand, there are no identities in the vector space Φ , that is

$$a_1 h_{t_1, x_1} + \dots + a_k h_{t_k, x_k} = 0 \Rightarrow a_1 = \dots = a_k = 0, \quad (2.21)$$

provided that $(k_i, x_i) \neq (k_j, x_j)$ for every $i \neq j$.

2.4 States on *-algebras and GNS representations

This short subsection might be skipped by the reader familiar with the notion of states on *-algebras and the GNS construction.

Quantization of the basic position and momentum variables for the scalar field amounts to finding a representation of the quantum algebra $\mathfrak{A}$ of the basic variables in the space of operators defined in a Hilbert space. In this paper we will be concerned with the so called GNS (Gel'fand-Naimark-Segal) representations defined by non-negative linear functional on the *-algebra $\mathfrak{A}$.

Recall, that:

Definition 2.8. *A state on a unital *-algebra $\mathfrak{A}$ is a non-trivial linear functional $\omega : \mathfrak{A} \rightarrow \mathbb{C}$, such that for every $\hat{a} \in \mathfrak{A}$*

$$\omega(\hat{a}^* \hat{a}) \geq 0, \quad \text{and} \quad \omega(\hat{1}) = 1, \quad (2.22)$$

where $\hat{1}$ stands for the unity element.

In fact, if we do not assume the equality above, it follows from the inequality (the positivity condition) itself, that

$$\omega(\hat{a}^*) = \overline{\omega(\hat{a})}, \quad \text{and} \quad \omega(\hat{1}) > 0. \quad (2.23)$$

Therefore, given the positivity, by rescaling we can always achieve the equality in Definition 2.8.

Given a state on $\mathfrak{A}$, we construct the corresponding GNS representation

$$(\mathcal{H}_\omega^0, \langle \cdot, \cdot \rangle_\omega, \rho_\omega, \Omega_\omega),$$

where: $\mathcal{H}_\omega^0$ is a vector space endowed with a unitary scalar product $\langle \cdot, \cdot \rangle_\omega$, ρ_ω a representation of $\mathfrak{A}$ on $\mathcal{H}_\omega^0$, and Ω_ω a vector in $\mathcal{H}_\omega^0$ which, when viewed as a state on $\mathfrak{A}$, coincides with ω (see Equation (2.30)). A detailed exposition of the GNS construction for algebras of unbounded operators can be found for example in [24]. Here we will only need the following elements and properties that are easy to prove:

(i) The vector space $\mathcal{H}_\omega^0$ is obtained as the quotient of $\mathfrak{A}$ considered as a vector space

$$\mathcal{H}_\omega^0 := \mathfrak{A}/J_\omega \quad (2.24)$$

where the vector subspace

$$J_\omega := \{ \hat{a} \in \mathfrak{A} : \omega(\hat{a}^* \hat{a}) = 0 \} \quad (2.25)$$

is a left ideal in $\mathfrak{A}$;

(ii) The unitary scalar product $\langle \cdot, \cdot \rangle_\omega$ is defined in $\mathcal{H}_\omega^0$ by the state ω , namely

$$\langle [\hat{a}], [\hat{b}] \rangle_\omega := \omega(\hat{a}^* \hat{b}), \quad (2.26)$$

where for every $\hat{c} \in \mathfrak{A}$, $[\hat{c}] \in \mathfrak{A}/J_\omega$ stands for the equivalence class defined by C ;

(iii) To every element $\hat{a}$ of $\mathfrak{A}$ we assign a linear (but in general unbounded) operator $\rho_\omega(\hat{a})$, defined in the entire vector space $\mathcal{H}_\omega^0$,

$$\rho_\omega(\hat{a}) : \mathcal{H}_\omega^0 \rightarrow \mathcal{H}_\omega^0 \quad (2.27)$$

$$\rho_\omega(\hat{a})[\hat{b}] := [\hat{a}\hat{b}]; \quad (2.28)$$

(iv) The vector Ω_ω is

$$\Omega_\omega = [\hat{1}], \quad (2.29)$$

and its relation with the state ω is

$$\omega(\hat{a}) = \langle \Omega_\omega, \rho_\omega(\hat{a})\Omega_\omega \rangle_\omega, \quad (2.30)$$

for every $\hat{a} \in \mathfrak{A}$;

(v) The map $\rho_\omega : \mathfrak{A} \rightarrow \text{End}(\mathcal{H}_\omega^0)$ is an homomorphism of the associative algebra $\mathfrak{A}$ into the associative algebra of the endomorphisms of the vector space $\mathcal{H}_\omega^0$.

(vi) The map ρ_ω is consistent with the *-structure of $\mathfrak{A}$ in the following way:

$$\langle \rho_\omega(\hat{a}^*)[\hat{b}], [\hat{c}] \rangle_\omega = \langle [\hat{b}], \rho_\omega(\hat{a})[\hat{c}] \rangle_\omega, \quad (2.31)$$

for every $\hat{a}, \hat{b}, \hat{c} \in \mathfrak{A}$.

A representation may be equivalent to a GNS representation. For the sake of precision we spell out the definition of equivalence:

Definition 2.9. Given $(V, (\cdot|\cdot), \rho, v_0)$, where: (a) V is a vector space endowed with a unitary form $(\cdot|\cdot)$, (b) ρ is an associative epimorphism

$$\rho : \mathfrak{A} \rightarrow \text{End}(V)$$

of a $*$ -algebra $\mathfrak{A}$ into the endomorphism algebra of V , and (c) $v_0 \in V$, we say, that $(V, (\cdot|\cdot), \rho, v_0)$ is equivalent to a given GNS representation $(\mathcal{H}_\omega^0, \langle \cdot, \cdot \rangle_\omega, \rho_\omega, \Omega_\omega)$ whenever there exists a unitary vector space isomorphism $\mathcal{I} : \mathcal{H}_\omega^0 \rightarrow V$, such that

$$\rho_\omega = \mathcal{I}^{-1} \circ \rho \circ \mathcal{I}, \quad \text{and} \quad \mathcal{I}(\Omega_\omega) = v_0.$$

Notice, that the notion of the Hilbert space is not necessary when one deals with the GNS algebras, as long as we are interested in the elements of the algebra $\mathfrak{A}$ only.

However, as for every unitary space, one can consider the Hilbert space

$$\mathcal{H}_\omega := \overline{\mathcal{H}_\omega^0}.$$

From this point of view, a difference between the current $*$ -algebra case and any other case of a C^* -algebra, is that since a general operator $\rho_\omega(\hat{a})$ is unbounded, an equality

$$\hat{a}^* = \hat{a} \tag{2.32}$$

if it takes place, does not imply that $\rho_\omega(\hat{a})$ acting in the domain $\mathcal{H}_\omega^0$ is essentially self-adjoint. If the algebra $\mathfrak{A}$ is commutative then, remarkably, (2.32) implies that the operator $\rho_\omega(\hat{a})$ admits a self-adjoint extension. Indeed, the operator commutes with the anti-linear operation

$$* : \mathcal{H}_\mu \rightarrow \mathcal{H}_\mu$$

induced by the $*$ -structure of $\mathfrak{A}$, in the commutative case. In general, the self-dual extension is not unique. However, due to the algebra of the problem, this unambiguity does not lead to any inconsistencies. Of course, given a $*$ -algebra, it is always interesting to ask which states define the GNS representations such that the real elements of the algebra correspond to the essentially-self adjoint operators.

2.5 Additional assumption about the representations of $\mathfrak{A}$

In this paper, we apply the GNS construction to the polymer $*$ -algebra $\mathfrak{A}$ (Definition 2.7).

Throughout this paper, we will be assuming about a state ω on $\mathfrak{A}$ that the corresponding GNS representation ρ_ω satisfies the following property:

Property 2.10. *For every point $x \in M$, the set of operators*

$$U_x = \{\rho_\omega(\hat{h}_{k,x}) : k \in \mathbb{R}\} \quad (2.33)$$

where $\hat{h}_{0,x}$ stands for the algebra unity element $\hat{1}$, is a group (with the operation of the composition), and

$$\rho_\omega(\hat{h}_{k,x})\rho_\omega(\hat{h}_{k',x}) = \rho_\omega(\hat{h}_{k+k',x}). \quad (2.34)$$

This assumption reflects the property of the exponentiated evaluation functions

$$e^{ik\phi(x)}e^{ik'\phi(x)} = e^{i(k+k')\phi(x)}. \quad (2.35)$$

It is analogous to the property

$$\rho_\omega(\hat{\pi}(f_1)) + \rho_\omega(\hat{\pi}(f_2)) = \rho_\omega(\hat{\pi}(f_1 + f_2)), \quad (2.36)$$

already satisfied as an identity due to the definition of the Lie algebra of the basic variables.

It follows from (2.34) that for every polymer position variable $h_{k,x}$, the operator

$$\rho_\omega(\hat{h}_{k,x}) : \mathcal{H}_\omega^0 \rightarrow \mathcal{H}_\omega^0 \quad (2.37)$$

is the unitary form preserving and bijective, therefore it is extendable by the continuity to a unitary operator in the Hilbert space $\mathcal{H}_\omega$. Indeed, it follows that:

$$\rho_\omega(\hat{h}_{-k,x})\rho_\omega(\hat{h}_{k,x}) = \rho(\hat{1}) = \text{id} = \rho_\omega(\hat{h}_{k,x})\rho_\omega(\hat{h}_{-k,x}), \quad (2.38)$$

$$\left\langle \rho_\omega(\hat{h}_{k,x})[\hat{a}], \rho_\omega(\hat{h}_{k,x})[\hat{b}] \right\rangle_\omega = \left\langle [\hat{a}], \rho_\omega(\hat{h}_{-k,x})\rho_\omega(\hat{h}_{k,x})[\hat{b}] \right\rangle_\omega = \left\langle [\hat{a}], [\hat{b}] \right\rangle. \quad (2.39)$$

The condition (2.34) is equivalent to the following condition imposed directly on the state ω

$$\omega(\hat{a}(\hat{h}_{k,y}\hat{h}_{k',y} - \hat{h}_{k+k',y})\hat{b}) = 0, \quad \text{for every } \hat{a}, \hat{b} \in \mathfrak{A}, \quad (2.40)$$

and every $k, k' \in \mathbb{R}, y \in M$.

We will be assuming the condition (2.34) in this paper. This is equivalent to considering states on the *-algebra

$$\tilde{\mathfrak{A}} = \mathfrak{A}/\tilde{J}, \quad (2.41)$$

where $\tilde{J}$ is the two-sided ideal generated by the set

$$\{\hat{h}_{k,y}\hat{h}_{k',y} - \hat{h}_{k+k',y} : y \in M, k, k' \in \mathbb{R}\}. \quad (2.42)$$

The canonical epimorphism

$$\mathfrak{A} \rightarrow \tilde{\mathfrak{A}}$$

is used to introduce the following notation for elements of $\tilde{\mathfrak{A}}$,

$$\begin{aligned}\hat{a} &\mapsto \tilde{a} \\ \hat{h}_{x,k} &\mapsto \tilde{h}_{x,k} \\ \hat{\pi}(f) &\mapsto \tilde{\pi}(f),\end{aligned}\tag{2.43}$$

in particular $\tilde{1}$ stands for the unity in $\tilde{\mathfrak{A}}$. A technical observation extensively used below is that every element of the quotient algebra $\tilde{\mathfrak{A}}$ can be represented by a finite linear combination of elements (2.19) such that $x_i \neq x_j$ for every pair $i \neq j$ and the hats are replaced by the tildes. In other words, for every finitely supported function $\mathbf{k} : M \rightarrow \mathbb{R}$ denote

$$\tilde{h}_{\mathbf{k}} = \prod_{x \in M} \tilde{h}_{x, \mathbf{k}(x)}.\tag{2.44}$$

The algebra $\tilde{\mathfrak{A}}$ is spanned by elements of the form

$$\tilde{h}_{\mathbf{k}}, \quad \text{and} \quad \tilde{h}_{\mathbf{k}} \prod_{j=1}^m \tilde{\pi}(f_j),\tag{2.45}$$

with all possible $\mathbf{k}$, $m \in \mathbb{N}$, and $f_1, \dots, f_m \in \mathcal{F}$.

Whereas, upon Property 2.10, each of the operators $\rho_{\omega}(\hat{h}_{k,x})$ will be unitary, operators $\rho_{\omega}(\hat{\pi}(f))$ may or may not be essentially self adjoint, depending on a state ω .

2.6 Symmetries

Suppose $\mathfrak{A}$ is the polymer $*$ -algebra corresponding to a momentum variable space $\Pi_{\mathcal{F}}$ (Definition 2.4) preserved by the group $\text{Diff}(M)$ of the diffeomorphisms of M (of a given class). Every diffeomorphism $\varphi \in \text{Diff}(M)$ acts on the pair representing the classical scalar field (ϕ, π) ,

$$\phi \mapsto (\varphi^{-1})^* \phi, \quad \pi \mapsto \varphi_* \pi,\tag{2.46}$$

(notice, that π , as a signed measure, is pushed forward by φ rather than pulled back by its inverse). The action passes naturally to the polymer Lie algebra and next to the polymer $*$ -algebra. φ defines a $*$ -automorphism of $\mathfrak{A}$

$$\sigma_{\varphi} : \mathfrak{A} \rightarrow \mathfrak{A}\tag{2.47}$$

$$\sigma_{\varphi}(\hat{h}_{k,x}) = \hat{h}_{k, \varphi(x)}\tag{2.48}$$

$$\sigma_{\varphi}(\hat{\pi}(f)) = \hat{\pi}((\varphi^{-1})^*(f)).\tag{2.49}$$

The automorphism σ_{φ} , as any automorphism of $\mathfrak{A}$, acts naturally in the space of the states defined on $\mathfrak{A}$. A state ω invariant with respect to that action of the diffeomorphism group $\text{Diff}(M)$ is called *diffeomorphism invariant*.

Moreover, consider an arbitrary *homeomorphism* φ . In non of the cases considered in this work the space of the smearing functions will be preserved by all the homeomorphisms. However, it may be true, that a given momentum variable $\pi(f) \in \Pi_{\mathcal{F}}$, the smearing function f is mapped by φ^{-1} into a function $(\varphi^{-1})^*f$ such that still $(\varphi^{-1})^*f \in \mathcal{F}$. The smearing functions of this property form a vector subspace of $\mathcal{F}$. Denote by Π_{φ} the corresponding vector subspace of Π . Explicitly,

$$\Pi_{\varphi} = \Pi_{\mathcal{F} \cap \varphi^* \mathcal{F}}$$

It comes with the polymer *-algebra, say $\mathfrak{A}_{\varphi}$, a subalgebra of $\mathfrak{A}$. It is obvious how to generalize the definition of the invariance of a state ω to the homeomorphisms. The formulae (2.48, 2.49) define a homomorphism

$$\sigma_{\varphi} : \mathfrak{A}_{\varphi} \rightarrow \mathfrak{A}. \quad (2.50)$$

Definition 2.11. *A state ω defined on the algebra $\mathfrak{A}$ is called homeomorphism invariant if for every homeomorphism φ*

$$\omega(\sigma_{\varphi}(a)) = \omega(a) \quad (2.51)$$

for every $a \in \mathfrak{A}_{\varphi}$.

Certainly, a homeomorphism invariant state on the polymer *-algebra $\mathfrak{A}$ is also diffeomorphism invariant.

For the completeness let us explain the relation between the diffeomorphism invariance of a state and the diffeomorphism covariance of the corresponding representation. A diffeomorphism invariant state ω on a polymer *-algebra $\mathfrak{A}$ defines a unitary representation U of the diffeomorphism group $\text{Diff}(M)$ in the unitary space $\mathcal{H}_{\omega}$ (2.24)

$$U(\varphi)[\hat{a}] = [\varphi^* \hat{a}]. \quad (2.52)$$

The representation ρ_{ω} is diffeomorphism covariant, in the sense

$$U(\varphi)\rho(\hat{a})U(\varphi)^{-1} = \rho(U(\varphi)\hat{a}) \quad (2.53)$$

for every $\hat{a} \in \mathfrak{A}$ and every $\varphi \in \text{Diff}(M)$.

In the next section, in the case of the momentum variable space $\Pi_{C_0^{m_0}(M, \mathbb{R})}$ we will show that a C^{m_0} diffeomorphism invariant state on $\mathfrak{A}$ is unique. This is the known one used in [21]. That state is also homeomorphism invariant.

In Sections 4 - 9 on the other hand, we will consider the momentum variable space $\Pi_{\mathcal{F}}$ spanned by characteristic functions of regions in M . In that case, we will additionally assume the homeomorphism invariance in order to derive a class of new states invariant with respect to the piecewise-analytic diffeomorphisms.

3 $\mathfrak{A}$ defined by $C_0^{m_0}(M, \mathbb{R})$ smearing functions

In this section we choose the space $\mathcal{F}$ of the smearing functions to be

$$\mathcal{F} = C_0^{m_0}(M, \mathbb{R}), \quad (3.1)$$

that is the space of the real C^{m_0} -functions of compact support and defined on M , where the differentiability class is fixed to be either an integer $m_0 \geq 0$, or $m_0 = \infty$, or $C^{(m_0)}$ stands for the semi-analytic functions according to [23].

We consider the polymer $*$ -star algebra $\mathfrak{A}$ corresponding to the momentum variable space $\Pi_{\mathcal{F}}$ defined by Definition 2.4 with the choice (3.1) of $\mathcal{F}$. We also identify $\Pi_{\mathcal{F}}$ and $\mathcal{F}$ via the vector space isomorphism (2.7).

In this section, the group $\text{Diff}(M)$ considered in Section 2.6 is the group $\text{Diff}^{m_0}(M)$ of all the C^{m_0} -diffeomorphisms of M . It turns out that the diffeomorphism invariance and the Property (2.10) determine a state ω on $\mathfrak{A}$ completely. That result, and the proof we present below are analogous the result of [23], but much simpler.

Theorem 3.1. *Suppose $\mathfrak{A}$ is the polymer $*$ -algebra (Definition 2.7) defined by $\mathcal{F} = C_0^{m_0}(M, \mathbb{R})$. On $\mathfrak{A}$, there exists exactly one C^{m_0} -diffeomorphism invariant state ω_0 such that the condition (2.40) (equivalent to Property 2.10) holds. The state is defined by*

$$\omega_0(\hat{a}\hat{\pi}(f)) = 0 = \omega_0(\hat{h}_{k_1, x_1} \dots \hat{h}_{k_n, x_n}), \quad (3.2)$$

for every $\hat{a} \in \mathfrak{A}$, every momentum variable $\pi(f) \in \Pi_{\mathcal{F}}$, and every set of labels $\{(k_1, x_1), \dots, (k_n, x_n)\}$ such that $x_i \neq x_j$ for every $i \neq j$, for all $i = 1, \dots, n$ and every $n \in \mathbb{N}$.

Proof. First, we show that if ω_0 is a diffeomorphism invariant state such Property (2.10) is true, then it satisfies (3.2). The main step is showing that

$$\omega_0(\hat{\pi}(f)^* \hat{\pi}(f)) = 0, \quad (3.3)$$

for every $f \in C_0^{m_0}(M, \mathbb{R})$. The trick is to notice that

Lemma 3.2. *For every $f \in C_0^{m_0}(M, \mathbb{R})$ there exists $g \in C_0^{m_0}(M, \mathbb{R})$ and $\epsilon > 0$ such that for every $-\epsilon < \lambda < \epsilon$, there is a $C_0^{m_0}(M, \mathbb{R})$ diffeomorphism $\phi_\lambda : M \rightarrow M$, such that*

$$g + \lambda f = \phi_\lambda^* g. \quad (3.4)$$

See [23] for the construction of ϕ_λ .

That is to say, $g + \lambda f$ is just a deformation of g , diffeomorphically trivial. Then the diffeomorphism invariance implies

$$\begin{aligned} \omega_0(\hat{\pi}(g)\hat{\pi}(g)) &= \omega_0(\hat{\pi}(g + \lambda f)\hat{\pi}(g + \lambda f)) \\ &= \omega_0(\hat{\pi}(g)\hat{\pi}(g)) + 2\lambda\omega_0(\hat{\pi}(f)\hat{\pi}(g)) + \lambda^2\omega_0(\hat{\pi}(f)\hat{\pi}(f)), \end{aligned} \quad (3.5)$$

for every value $\lambda \in]-\epsilon, \epsilon[$. Hence, the terms proportional to λ and λ^2 vanish. (Recall also that $\hat{\pi}(f)^* = \hat{\pi}(f)$.) This implies the first equality in (3.2). Notice, that it also implies

$$\omega_0(\hat{\pi}(f)A) = \overline{\omega_0(A^*\hat{\pi}(f))} = 0, \quad (3.6)$$

for every $A \in \mathfrak{A}$ and every $\pi(f) \in \Pi_{\mathcal{F}}$.

Now, consider an element $\hat{h} := \hat{h}_{k_1, x_1} \dots \hat{h}_{k_n, x_n} \in \mathfrak{A}$. Certainly, for every $f \in C_0^{m_0}(M, \mathbb{R})$

$$0 = \omega_0(\hat{h}h\hat{\pi}(f) - \hat{\pi}(f)h) = \left(\sum_{i=1}^n k_i f(x_i) \right) \omega_0(h). \quad (3.7)$$

If $\hat{h}$ satisfies the assumption that $x_i \neq x_j$ whenever $i \neq j$ then there is $f \in C_0^{m_0}(M, \mathbb{R})$ such that the coefficient at $\omega_0(\hat{h})$ in the equality above is not zero, hence the second equality in (3.2) is necessarily true. Hence, we have showed that (3.2) is a necessary condition.

Now, we show, that if ω_0 satisfies (Property 2.10) and (3.2), then it is unique. Notice, that the only elements of $\mathfrak{A}$ that have not been taken into account in (3.2) are the linear combinations of elements $\hat{h}_{k_1, y_1} \dots \hat{h}_{k_n, y_n}$ such that there is $i, j \in \{1, \dots, k\}$ such that

$$i \neq j, \quad \text{and} \quad y_i = y_j = x. \quad (3.8)$$

However, the equations (3.2, 2.40) and the second equation in (2.8) determine ω_0 even on those elements of $\mathfrak{A}$.

Concluding, we have shown that if ω_0 is a diffeomorphism invariant state such that Property (2.10) is satisfied, then it satisfies (3.2) and is unique.

It is easy to construct a state ω_0 by using (2.34, 3.2). The conditions (2.34, 3.2) define a state on the algebra $\tilde{\mathfrak{A}}$ (see 2.41). The state ω_0 is the pullback of that state to $\mathfrak{A}$, hence it exists. Obviously, it is diffeomorphism invariant. $\square$

The GNS representation ρ_{ω_0} defined by the state ω_0 can be derived explicitly. The first step is to find a convenient notation for the elements of the quotient vector space $\mathcal{H}_{\omega_0}^0$.

Every non-zero element of $[\hat{a}] \in \mathfrak{A}/J_{\omega_0}$ can be defined by $\hat{a} \in \mathfrak{A}$ of the following form,

$$\hat{a} = \left[\prod_{i=1}^n \hat{h}_{k_i, x_i} \right]$$

and can be labeled by a finitely supported function $\mathbf{k} : M \rightarrow \mathbb{R}$,

$$\mathbf{k} = \sum_{i=1}^n k_i \mathbf{1}_{\{x_i\}}, \quad (3.9)$$

where given $x \in M$, $\mathbf{1}_{\{x\}}$ is the characteristic function of x . In particular, the identically zero function labels the element $[\hat{1}] \in \mathfrak{A}/J_{\omega_0}$. Therefore we will write

$$|\mathbf{k}\rangle := [\prod_{x \in M}^n \hat{h}_{\mathbf{k}(x),x}] \quad (3.10)$$

where $\mathbf{k} : M \rightarrow \mathbb{R}$ has a finite support.

Therefore, $\mathcal{H}_{\omega_0}^0$ is in this case the complex vector space Cyl of the finite formal linear combinations of elements $|\mathbf{k}\rangle$ labeled by all the functions $\mathbf{k} : M \rightarrow \mathbb{R}$ of *finite* support,

$$\mathcal{H}_{\omega_0}^0 = \text{Cyl} = \left\{ \sum_{i=1}^n a_i |\mathbf{k}_i\rangle : n \in \mathbb{N}, a_i \in \mathbb{C}, \mathbf{k}_i : M \rightarrow \mathbb{R} \text{ finitely supported} \right\}. \quad (3.11)$$

Now, the GNS unitary scalar product $\langle \cdot, \cdot \rangle_{\omega_0}$ defined in $\mathcal{H}_{\omega_0}^0$ by the state ω_0 is the following unitary scalar product $(\cdot | \cdot)_{\text{Cyl}}$:

$$(|\mathbf{k}_1\rangle | |\mathbf{k}_2\rangle)_{\text{Cyl}} = \begin{cases} 1, & \text{if } \mathbf{k}_1 = \mathbf{k}_2 \\ 0, & \text{if } \mathbf{k}_1 \neq \mathbf{k}_2. \end{cases} \quad (3.12)$$

Finally, the action $\rho_{\omega_0} : \mathfrak{A} \rightarrow \text{End}(\text{Cyl})$ defined on the generators of $\mathfrak{A}$ and the basis in Cyl can be derived:

$$\rho_{\omega_0}(\hat{h}_{k',x})(|\mathbf{k}\rangle) = [\hat{h}_{k',x} \prod_{x \in M}^n \hat{h}_{\mathbf{k}(x),x}] = |\mathbf{k} + k' \mathbf{1}_{\{x\}}\rangle \quad (3.13)$$

$$\begin{aligned} \rho(\hat{\pi}(f))(|\mathbf{k}\rangle) &= [\hat{\pi}(f) \prod_{x \in M}^n \hat{h}_{\mathbf{k}(x),x}] = [[\hat{\pi}(f), \prod_{x \in M}^n \hat{h}_{\mathbf{k}(x),x}]] = \\ &= \sum_{x \in M} \mathbf{k}(x) f(x) |\mathbf{k}\rangle. \end{aligned} \quad (3.14)$$

where the product (sum) has only finitely many non-unit (non-zero) factors (terms).

Some more remarks are in order. That unique state is used in LQG for the quantum scalar field coupled with the quantum geometry [20, 21]. The symmetry group of the quantum geometry defined by LQG is the group of semi-analytic diffeomorphisms. As it was shown in [23], Lemma 3.2 continues to be true if the C^{m_0} differentiability class is replaced by the semi-analyticity assumption defined therein. Therefore the theorem is true in that case as well.

The representation ρ_{ω_0} is referred to in the literature, as the polymer representation. For the same polymer $\ast$ -algebra $\mathfrak{A}$ one may also define the standard Fock state of the scalar field. The comparison between the polymer representation and the Fock representation was discussed in [21].

An important subtlety of Theorem 3.1 is that both: the smearing functions used in the definition of the basic momentum variables, and the diffeomorphisms $M \rightarrow M$ are assumed to be of the same class C^{m_0} . The reason is that in the proof of Lemma 3.2 (see[23]), the suitable diffeomorphism is constructed from a given smearing function. However, if we fix the differentiability class of the diffeomorphism-symmetry group as C^{m_0} , but consider the smearing functions of the class $C^{m'}$ with $m' < m_0$, the proof fails, and, in fact, the theorem is not true. (It is easy to construct a counter example by using the counter example to the uniqueness concerning the quantum algebra of the holonomy-flux observable found in [23]).

In the current work, our aim is to use that observation in investigation of the issue of the existence of new diffeomorphism invariant states.

A natural alternative to the $C_0^{m_0}$ condition used in Theorem 3.1 is smearing the momentum variables against the characteristic functions of subsets in M . (This also one of the choices typically used in LQG [20]). With this choice of the space $\mathcal{F}$ in Definition 2.4, we will be able to construct a family of new states.

4 The CW-complex *-algebra

4.1 The CW-complex vector space $\mathcal{F}_S$

4.1.1 Definitions, the Euler Characteristics

We assume in this section that the manifold M is endowed with a piecewise-analytic structure, and consider piecewise-analytic k -simplexes in M , both notions precisely defined in Appendix. In particular, if S is a k -simplex in M , then, depending on k , S is: (0) a 1-point subset of M if $k = 0$, (i) an ‘open’ interval embedded in M if $k = 1$, (N) an open ball in $\mathbb{R}^{\dim M}$ embedded in M with an inverse local chard, if the $k = \dim M$. In the last case S is referred to just as a ball in M . In the general case S is referred to just as simplex (we skip ‘piecewise-analytic’).

Definition 4.1. Denote by $\mathcal{S}$ the set of all the k -simplexes in M , where k ranges from 0 to $\dim M$.

Given a subset $U \in M$, by the characteristic function $\mathbf{1}_U$ we mean the function

$$\mathbf{1}_U : M \rightarrow \mathbb{R} \tag{4.1}$$

$$\mathbf{1}_U(x) = \begin{cases} 1, & \text{if } x \in U \\ 0, & \text{if } x \notin U. \end{cases} \tag{4.2}$$

Definition 4.2. The CW-complex vector space $\mathcal{F}_S$ is the real vector space spanned by

the linear combinations of the characteristic functions $\mathbf{1}_S$ of all the k -simplexes S in M :

$$\mathcal{F}_S = \left\{ \sum_{i=1}^n a_i \mathbf{1}_{S_i} : n \in \mathbb{N}, a_i \in \mathbb{R}, S_i \in \mathcal{S} \right\}.$$

The key identities satisfied by the characteristic functions valid for arbitrary subsets $U_1, U_2 \subset M$ are

$$\mathbf{1}_{U_1} + \mathbf{1}_{U_2} - \mathbf{1}_{U_1 \cap U_2} - \mathbf{1}_{U_1 \cup U_2} = 0 \quad (4.3)$$

$$\mathbf{1}_{U_1 \setminus U_2} - \mathbf{1}_{U_1} + \mathbf{1}_{U_1 \cap U_2} = 0. \quad (4.4)$$

Due to them, despite of the fact, we have used only the simplexes to define the space $\mathcal{F}_S$, there are many other subsets $U \subset M$, such that the characteristic function

$$\mathbf{1}_U \in \mathcal{F}_S.$$

Indeed, this is the case whenever U is triangulable (see Appendix).

As a warm up, and to introduce the Euler Characteristic, consider a linear map

$$\chi : \mathcal{F}_S \rightarrow \mathbb{R} \quad (4.5)$$

and assume it is homeomorphism invariant in the sense, that the number $\chi(\mathbf{1}_S)$ depends only on the dimension of the simplex S . It is easy to see that this map is uniquely determined by the value of $\chi(S_0)$ where S_0 is a 0-simplex, meaning a one point subset of M . Indeed, if S_1 is a 1-simplex, then we can always split it into a disjoint union of two 1-simplexes and one 0-simplex, and consider the corresponding identity between the characteristic functions

$$\begin{aligned} S_1 &= S'_1 \cup S'_0 \cup S''_1 \\ \mathbf{1}_{S_1} &= \mathbf{1}_{S'_1} + \mathbf{1}_{S'_0} + \mathbf{1}_{S''_1}. \end{aligned} \quad (4.6)$$

The linearity and the homeomorphism invariance of χ applied to the identity imply

$$\chi(\mathbf{1}_{S_1}) = -\chi(\mathbf{1}_{S'_0}).$$

Repeating exactly the same calculation for 2-simplex up to $\dim M$ -simplex, one can convince himself, that for every k -simplex S_k ,

$$\chi(\mathbf{1}_{S_k}) = \mu_0 (-1)^k, \quad (4.7)$$

where μ_0 is an arbitrary constant. The map

$$\mathcal{S} \ni S_k \mapsto \chi_E(S_k) = (-1)^k \in \mathbb{R}, \quad (4.8)$$

where k is the dimension of S_k , is called the Euler Characteristic.

Thus far we have derived a necessary condition determining the possibly existing linear map $\chi : \mathcal{F}_S \rightarrow \mathbb{R}$ up to the constant μ_0 . Another argument is needed to show, that the linear map χ exists at all. The existence relies on two facts. The first one, is that for every n -tuple of the simplexes $S_1, \dots, S_n \in \mathcal{S}$, there is a single triangulation, that is a finite subset $\mathcal{S}' \subset \mathcal{S}$ of pairwise disjoint simplexes in M such that

$$S_i = \bigcup_{j=1}^{n_i} S'_{ij}, \quad S'_{ij} \in \mathcal{S}'$$

for every $i = 1, \dots, n$, and $j = 1, \dots, n_i$. The second fact is, that the Euler Characteristic is triangulation independent, that is for every simplex S and its triangulation, that is a decomposition

$$S = \bigcup_{i=1}^n S_i$$

into the union of pairwise disjoint simplexes $S_1, \dots, S_n$, the identity holds

$$\chi_E(S) = \sum_{i=1}^n \chi_E(S_i).$$

We hope it will not lead to any confusion if we denote by

$$\chi_E : \mathcal{F}_S \rightarrow \mathcal{F}_S$$

the linear extension of the Euler Characteristic, that is the map (4.8) normalized by the condition

$$\mu_0 = 1,$$

for every 0-simplex in M . We keep calling the extension $\chi_E : \mathcal{F}_S \rightarrow \mathbb{R}$ the Euler Characteristic functional.

4.1.2 Homeomorphism invariant, symmetric, bilinear forms

On the CW-complex vector space $\mathcal{F}_S$, consider a symmetric, bilinear form

$$(\cdot|\cdot) : \mathcal{F}_S \times \mathcal{F}_S \rightarrow \mathbb{R},$$

non-negative, that is such that for every $s \in \mathcal{F}_S$

$$(s|s) \geq 0.$$

We also assume, that the form assigns to a pair of simplexes $S_1, S_2 \in \mathcal{S}$ a number $(\mathbf{1}_{S_1}|\mathbf{1}_{S_2})$ depending only on the homeomorphic characteristics of the pair $\{S_1, S_2\}$. Precisely, we assume that $(\cdot|\cdot)$ is homeomorphism invariant in the following sense:

Definition 4.3. A bilinear form $(\cdot|\cdot) : \mathcal{F}_S \times \mathcal{F}_S \rightarrow \mathbb{R}$ is called homeomorphism invariant if for every pair of simplexes $S_1, S_2 \in \mathcal{S}$ and every homeomorphism $\phi : M \rightarrow M$ such that $\phi(S_1), \phi(S_2) \in \mathcal{S}$, the form satisfies

$$(\mathbf{1}_{\phi(S_1)} | \mathbf{1}_{\phi(S_2)}) = (\mathbf{1}_{S_1} | \mathbf{1}_{S_2}). \quad (4.9)$$

In this section we will find all the homeomorphism invariant, bilinear, symmetric, non-negative forms on the CW-complex vector space $\mathcal{F}_S$.

The key step is the following Lemma:

Proposition 4.4. Suppose that, $\dim M > 1$ and M is connected. Suppose $(\cdot|\cdot) : \mathcal{F}_S \times \mathcal{F}_S \rightarrow \mathbb{R}$ is a homeomorphism invariant symmetric, bilinear, non-negative form. Then, for any two balls $B_0, B_1 \in \mathcal{S}$

$$(\mathbf{1}_{B_1} - \mathbf{1}_{B_0} | \mathbf{1}_{B_1} - \mathbf{1}_{B_0}) = 0. \quad (4.10)$$

Proof. Consider first the special case of the balls, when $B_0 \subset B_1$ and the difference $B_1 \setminus B_0$ is a tube¹, that is, there is a local coordinate system $(x^1, \dots, x^N)$ in M such that

$$B_1 \setminus B_0 = R = \{ (x^1, \dots, x^N) \mid r_0^2 \leq (x^1)^2 + (x^2)^2 < r_1^2, |x_i| < H, 2 \leq i \leq N \}. \quad (4.11)$$

We have,

$$\mathbf{1}_{B_1} - \mathbf{1}_{B_0} = \mathbf{1}_R,$$

therefore, in particular $\mathbf{1}_R \in \mathcal{F}_S$.

Lemma 4.5. Suppose $R \subset M$ is a tube defined in a local coordinate system by (4.11). Suppose M and $(\cdot|\cdot)$ satisfy the assumptions of Proposition 4.4. Then,

$$(\mathbf{1}_R | \mathbf{1}_R) = 0.$$

Proof. The trick consists in cutting out the tube R into n disjoint, homeomorphic pieces

$$R = R_1^{(n)} \cup \dots \cup R_n^{(n)},$$

such that each two pairs of pairs $(R, R_I^{(n)})$ and $(R, R_J^{(n)})$ are homeomorphic to each other and

$$\mathbf{1}_{R_I^{(n)}} \in \mathcal{S}$$

for every $I = 1, \dots, n$. Indeed, we can choose some $n \in \mathbb{N}$ and using the same coordinates define

$$R_I^{(n)} = \{ (x^1, \dots, x^N) \in R : x^1 = r \cos \phi, x^2 = r \sin \phi, r \in \mathbb{R}, (I-1)\frac{2\pi}{n} \leq \phi < I\frac{2\pi}{n} \}$$

¹It is useful to have in mind the identification of a ball with a cylinder

Then,

$$\mathbf{1}_R = \mathbf{1}_{R_1^{(n)}} + \cdots + \mathbf{1}_{R_n^{(n)}}, \text{ and } \mathbf{1}_{R_I^{(n)}} \in \mathcal{F}_S$$

for every $I = 1, \dots, n$. Now, the number in question can be written in the following form,

$$(\mathbf{1}_R | \mathbf{1}_R) = n(\mathbf{1}_R | \mathbf{1}_{R_1^{(n)}}) \quad (4.12)$$

Where we have used the homeomorphism invariance of the form $(\cdot | \cdot)$. The left hand side of the equality is independent of n . The right hand side obviously depends on n . However, the factor $(\mathbf{1}_R | \mathbf{1}_{R_1^{(n)}})$, again due to the homeomorphism invariance, is n independent! The only solution of the equation (4.12) valid for arbitrary n is

$$(\mathbf{1}_R | \mathbf{1}_R) = 0.$$

□

In the continuation of the proof of Proposition 4.4 we will use the known fact, that given in a vector space a symmetric, bi-linear and non-negative form, the ‘zero norm’ vectors form a vector space, that is, in our case

$$(s | s) = 0, \text{ and } (s' | s') = 0 \Rightarrow (s + as' | s + as') = 0$$

for every $a \in \mathbb{R}$.

Lemma 4.6. *Suppose B_0 and B'_0 are two disjoint balls in M (that is $B_0, B'_0 \in \S$ are simplexes of the maximal dimension). Suppose M and $(\cdot | \cdot)$ satisfy the assumptions of Proposition 4.4. Then,*

$$(\mathbf{1}_{B_0} - \mathbf{1}_{B'_0} | \mathbf{1}_{B_0} - \mathbf{1}_{B'_0}) = 0.$$

Proof. Given the balls B_0 and B'_0 , there is a single ball B_1 such that both (i) $B_0 \subset B_1$ and $R = B_1 \setminus B_0$ is a tube, and (ii) $B'_0 \subset B_1$ and $R' = B_1 \setminus B'_0$ is a tube. (The notion of ‘tube’ is defined by (4.11)). We can write

$$\mathbf{1}_{B_0} - \mathbf{1}_{B'_0} = \mathbf{1}_{B_0} - \mathbf{1}_{B_1} + \mathbf{1}_{B_1} - \mathbf{1}_{B'_0} = -\mathbf{1}_R + \mathbf{1}_{R'}.$$

The application of Lemma 4.5 concludes the proof of Lemma 4.6. □

Finally, suppose $B_1, B_2 \subset M$ are arbitrary two balls in M . Let B_3 be a ball in M disjoint from $B_1 \cup B_2$. We have

$$\mathbf{1}_{B_1} - \mathbf{1}_{B_2} = \mathbf{1}_{B_1} - \mathbf{1}_{B_3} + \mathbf{1}_{B_3} - \mathbf{1}_{B_2}$$

and Lemma 4.6 applies to the first difference as well as to the second one. Hence, Proposition 4.4 follows. □

The conclusion from Proposition 4.4, is that that if we fix a ball B_0 in M , then for every other ball B and every $s \in \mathcal{F}_S$,

$$(\mathbf{1}_B|s) = (\mathbf{1}_{B_0}|s)$$

provided the assumptions of Proposition 4.4 are satisfied.

It follows, that the form $(\cdot|\cdot)$ can be determined via Proposition 4.4 up a normalization factor. To see it, note first:

Lemma 4.7. *For every simplex S in M , there exist balls $B_1, \dots, B_k$, such that*

$$\mathbf{1}_S = \sum_{i=1}^n m_i \mathbf{1}_{B_i}, \quad (4.13)$$

where $m_1, \dots, m_n$ are integers such that

$$\chi_E(S) = (-1)^{\dim M} \sum_{i=1}^n m_i. \quad (4.14)$$

Proof. The formula (4.14) follows directly from the application of the Euler Characteristic functional to (4.13). Given a k -simplex S in M such that $k < \dim M$, it is easy to see that there are three $k+1$ simplexes S', S'' and S''' in M such that

$$\mathbf{1}_S = \mathbf{1}_{S'''} - \mathbf{1}_{S''} - \mathbf{1}_{S'}$$

and all the terms on the right hand-side are characteristic functions of $k+1$ -simplexes. Repeating this observation sufficiently many times we get (4.13). $\square$

Therefore, we can calculate now for arbitrary two elements $s, s' \in \mathcal{F}_S$

$$(s|s') = \left(\sum_{i=1}^n m_i \mathbf{1}_{B_i} \middle| \sum_{i'=1}^{n'} m_{i'} \mathbf{1}_{B_{i'}} \right) = \chi_E(s) \chi_E(s') (\mathbf{1}_{B_0} | \mathbf{1}_{B_0}), \quad (4.15)$$

where we have used that

$$(\mathbf{1}_{B_i} | \mathbf{1}_{B_{i'}}) = (\mathbf{1}_{B_0} | \mathbf{1}_{B_0}).$$

Hence, $(\cdot|\cdot)$ is determined indeed up to a constant factor. Due to the independence of the Euler Characteristic of triangulation, the formula above does define a bilinear form $(\cdot|\cdot) : \mathcal{F}_S \times \mathcal{F}_S \rightarrow \mathbb{R}$, given any value

$$\lambda = (\mathbf{1}_{B_0} | \mathbf{1}_{B_0})$$

In this way we have proved, that:

Theorem 4.8. *There exist a unique, modulo the re-scaling by an arbitrary factor $\mu_2 > 0$, homeomorphism invariant, bilinear, symmetric form $(\cdot|\cdot)$ defined on the CW-complex vector space $\mathcal{F}_S$, provided $\dim M > 1$ and M is connected. The form is defined as follows:*

$$(\cdot|\cdot) = \mu_2 \chi_E(\cdot) \chi_E(\cdot), \quad (4.16)$$

where $\chi_E : \mathcal{F}_S \rightarrow \mathbb{R}$ is the Euler Characteristic functional .

Notice, finally, that it follows from Theorem 4.8 that the resulting unitary space

$$\mathcal{F}_S / \{s \in \mathcal{F}_S : (s|s) = 0\}$$

is just one dimensional. If we denote by $[\mathbf{1}_B]$ the element of the space $\mathcal{F}_S / \{s \in \mathcal{F}_S : (s|s) = 0\}$ corresponding to a ball B in M , then the projection map is

$$\mathcal{F}_S \ni s \mapsto (-1)^{\dim M} \chi_E(s) [\mathbf{1}_B].$$

Notice also, that the element $[\mathbf{1}_B]$ is unique, it does not depend on which ball B in M we use to define it.

4.2 Homeomorphism invariant states on $\exp(\odot \mathcal{F}_S^{\mathbb{C}})$

Consider the complexification $\mathcal{F}_S^{\mathbb{C}}$ of the (real) CW-complex vector space $\mathcal{F}_S$. We will denote by $\odot$ the symmetrized tensor product, and use the following notation for an element $s \in \mathcal{F}_S^{\mathbb{C}}$

$$s^{\odot n} = s \odot \cdots \odot s \quad (4.17)$$

(n copies of s on the right hand side) and extend this notation to the space $\mathcal{F}_S^{\mathbb{C}}$ itself

$$(\mathcal{F}_S^{\mathbb{C}})^{\odot n} = \mathcal{F}_S^{\mathbb{C}} \odot \cdots \odot \mathcal{F}_S^{\mathbb{C}}, \quad (4.18)$$

that is $(\mathcal{F}_S^{\mathbb{C}})^{\odot n}$ is the space of the symmetric elements of the tensor product $\mathcal{F}_S^{\mathbb{C}} \otimes \cdots \otimes \mathcal{F}_S^{\mathbb{C}}$. Consider the following vector space

$$\exp(\odot \mathcal{F}_S^{\mathbb{C}}) := \bigoplus_{n=0}^{\infty} (\mathcal{F}_S^{\mathbb{C}})^{\odot n}. \quad (4.19)$$

The space has the natural commutative $*$ -algebra structure. Indeed, $\exp(\odot \mathcal{F}_S^{\mathbb{C}})$ is by definition a complex vector space. The associative composition operation is the symmetrized tensor product $\odot$, and the star operation is defined by using the complex conjugation $\bar{\cdot}$ in $\mathcal{F}_S^{\mathbb{C}}$,

$$\begin{aligned} (a \odot \cdots \odot b)(c \odot \cdots \odot d) &= a \odot \cdots \odot b \odot c \odot \cdots \odot d \\ (a \odot \cdots \odot b)* &= \bar{b} \odot \cdots \odot \bar{a}. \end{aligned} \quad (4.20)$$

Definition 4.9. The $*$ -algebra $\exp(\odot \mathcal{F}_{\mathcal{S}}^{\mathbb{C}})$ is called the CW-complex $*$ -algebra.

As we will see this algebra is naturally isomorphic to subalgebra of $\mathfrak{A}$ generated by momentum variables 5.1.

Definition 4.10. A state $\check{\omega} : \exp(\odot \mathcal{F}_{\mathcal{S}}^{\mathbb{C}}) \rightarrow \mathbb{C}$ is called homeomorphism invariant if for every k -tuple $S_1, \dots, S_k \in \mathcal{S}$ of simplexes in M and every homeomorphism $\varphi : M \rightarrow M$ such that still $\varphi(S_1), \dots, \varphi(S_k) \in \mathcal{S}$, the following is true

$$\check{\omega}(\mathbf{1}_{\varphi(S_1)} \odot \dots \odot \mathbf{1}_{\varphi(S_k)}) = \check{\omega}(\mathbf{1}_{S_1} \odot \dots \odot \mathbf{1}_{S_k}).$$

In this section we derive all the homeomorphism invariant states on $\exp(\odot \mathcal{F}_{\mathcal{S}}^{\mathbb{C}})$.

Suppose that

$$\check{\omega} : \exp(\odot \mathcal{F}_{\mathcal{S}}^{\mathbb{C}}) \rightarrow \mathbb{C}$$

is a homeomorphism invariant state. It is clear that $\check{\omega}$ defines a homeomorphism invariant, symmetric, bilinear form $(\cdot | \cdot) : \mathcal{F}_{\mathcal{S}} \times \mathcal{F}_{\mathcal{S}} \rightarrow \mathbb{R}$, namely

$$(s | s') = \check{\omega}(s \odot s')$$

(recall, that the subspace $\mathcal{F}_{\mathcal{S}} \subset \mathcal{F}_{\mathcal{S}}^{\mathbb{C}}$ consist of the real elements such that $\bar{s} = s$). Given any two balls B_1 and B_2 in M , Proposition 4.4 guarantees that the difference $\mathbf{1}_{B_1} - \mathbf{1}_{B_2} \in \mathcal{F}_{\mathcal{S}}$ satisfies

$$\check{\omega}(\mathbf{1}_{B_1} - \mathbf{1}_{B_2}) \odot (\mathbf{1}_{B_1} - \mathbf{1}_{B_2}) = 0.$$

Now, it follows from the basic properties of the commutative $*$ -algebras, that for every $A, A' \in \exp(\odot \mathcal{F}_{\mathcal{S}}^{\mathbb{C}})$,

$$\check{\omega}(A \odot (\mathbf{1}_{B_1} - \mathbf{1}_{B_2}) \odot A') = 0. \quad (4.21)$$

Using this observation, we conclude that the state $\check{\omega}$ is completely characterized by the following sequence of the numbers

$$\mu_n := \check{\omega}(\mathbf{1}_{B_0}^{\odot n}), \quad n \in \mathbb{N}. \quad (4.22)$$

where $B_0 \in \mathcal{S}$ is an arbitrarily fixed ball in M . Indeed, given any finite set of balls $B_1, \dots, B_n$ in M the identity (4.21) applied n times implies

$$\check{\omega}(\mathbf{1}_{B_1} \odot \dots \odot \mathbf{1}_{B_n}) = \check{\omega}((\mathbf{1}_{B_0})^{\odot n}).$$

Furthermore, due to (4.13, 4.14), the subset of all the elements of $\exp(\odot \mathcal{F}_{\mathcal{S}}^{\mathbb{C}})$ of the form $\mathbf{1}_{B_1} \odot \dots \odot \mathbf{1}_{B_n}$ where B_k are balls in M , spans the vector space $\exp(\odot \mathcal{F}_{\mathcal{S}}^{\mathbb{C}})$.

In other words, every homeomorphism invariant state $\check{\omega}$ on $\exp(\odot \mathcal{F}_{\mathcal{S}}^{\mathbb{C}})$ is unambiguously determined by its restriction to the unital subalgebra $\mathbb{C}[\mathbf{1}_{B_0}]$ of $\exp(\odot \mathcal{F}_{\mathcal{S}}^{\mathbb{C}})$

generated by an element $\mathbf{1}_{B_0}$, where B_0 is an arbitrarily fixed ball in M . It is clear that $\mathbb{C}[\mathbf{1}_{B_0}]$ is isomorphic to the $*$ -algebra $\mathbb{C}[\tau]$ of complex valued polynomials of one real variable; let us explain the notation:

$$\tau : \mathbb{R} \rightarrow \mathbb{R}, \quad \tau(r) = r$$

is the identity map,

$$\mathbb{C}[\tau] = \left\{ \sum_{j=1}^n a_j \tau^{j-1} : n \in \mathbb{N}, a_j \in \mathbb{C} \right\}.$$

and the isomorphism $\mathbb{C}[\mathbf{1}_{B_0}] \rightarrow \mathbb{C}[\tau]$ is defined just by

$$\mathbf{1}_{B_0} \mapsto \tau, . \tag{4.23}$$

Thus we have arrived at

Lemma 4.11. *Every homeomorphism invariant state $\tilde{\omega}$ on the $*$ -algebra $\exp(\odot \mathcal{F}_{\mathcal{S}}^{\mathbb{C}})$ is determined by a state μ defined on the $*$ -algebra $\mathbb{C}[\tau]$, according to the following formula*

$$\tilde{\omega}(\mathbf{1}_{B_1} \dots \mathbf{1}_{B_k}) = \mu(\tau^k), \tag{4.24}$$

for every $k \in \mathbb{N}$.

In Lemma 4.11 the existence of the state $\tilde{\omega}$ defined on $\exp(\odot \mathcal{F}_{\mathcal{S}}^{\mathbb{C}})$ has been assumed. Now, we turn to the existence issue itself. We will show, that:

Lemma 4.12. *Every state μ defined on the polynomial $*$ -algebra $\mathbb{C}[\tau]$ defines a unique homeomorphism invariant state $\tilde{\omega}$ on $\exp(\odot \mathcal{F}_{\mathcal{S}}^{\mathbb{C}})$ such that ((4.24)).*

Proof. Let us begin by constructing a $*$ -homomorphism

$$\text{Eul} : \exp(\odot \mathcal{F}_{\mathcal{S}}^{\mathbb{C}}) \rightarrow \mathbb{C}[\tau]$$

such that for every ball B' in M ,

$$\mathbf{1}_{B'} \mapsto \tau \in \mathbb{C}[\tau].$$

The linear extension of the equality above to the vector space $\mathcal{F}_{\mathcal{S}}^{\mathbb{C}}$, the complexification of $\mathcal{F}_{\mathcal{S}}$, is defined by using the Euler Characteristic (4.8) in the following way,

$$\begin{aligned} \text{Eul} : \mathcal{F}_{\mathcal{S}}^{\mathbb{C}} &\rightarrow \mathbb{C}[\tau], \\ s &\mapsto (-1)^{\dim M} \chi_E(s) \tau \end{aligned} \tag{4.25}$$

Because $\mathcal{F}_S^{\mathbb{C}}$ generates the commutative algebra $\exp(\odot \mathcal{F}_S^{\mathbb{C}})$ freely modulo the linear relations satisfied by its elements, the above formula can be uniquely extended to the whole algebra $\exp(\odot \mathcal{F}_S^{\mathbb{C}})$,

$$\text{Eul}(s_1 \odot \cdots \odot s_n) := \text{Eul}(s_1) \cdots \text{Eul}(s_n) \in \mathbb{C}[\tau], \quad (4.26)$$

Suppose μ is a state on $\mathbb{C}[\tau]$. The pullback

$$\tilde{\omega} := \mu \circ \text{Eul}$$

is certainly a state on $\exp(\odot \mathcal{F}_S^{\mathbb{C}})$. It is also homeomorphism invariant. Finally, the state $\tilde{\omega}$ satisfies ((4.24)). □

Let us summarize our observations by the following:

Theorem 4.13. *There is a 1-1 correspondence between the space of the homeomorphism invariant states on the CW-complex *-algebra $\exp(\odot \mathcal{F}_S^{\mathbb{C}})$ and the space of the states on the polynomial algebra $\mathbb{C}[\tau]$. The correspondence is defined by the pullback Eul^* with the *-homomorphism $\text{Eul} : \exp(\odot \mathcal{F}_S^{\mathbb{C}}) \rightarrow \mathbb{C}[\tau]$ (4.25, 4.26).*

The careful reader noticed, that the polynomial algebra $\mathbb{C}[\tau]$ has emerged just as isomorphic to the unital subalgebra of the CW-complex algebra, generated by a fixed single ball $\mathbf{1}_{B_0}$. In fact, using the polynomial algebra representation has many advantages. To begin with, an example of a state μ on the polynomial algebra $\mathbb{C}[\tau]$ is defined by a probability (regular, Borel) measure $d\mu$ on $\mathbb{R}$, via

$$\mu(\tau^k) := \int_{\mathbb{R}} \tau^k d\mu(\tau), \quad (4.27)$$

for every $k \in \mathbb{N}$, provided all the functions τ^k are integrable. Conversely, every state μ on the polynomial *-algebra $\mathbb{C}[\tau]$ can be constructed in that way. This nontrivial fact follows from the existence of a self-adjoint extension for the operator $\rho_\mu(\tau)$, where ρ_μ is the GNS representation (see Section 2.4. The measure is unique if the extension is unique).

Given a homeomorphism invariant state $\tilde{\omega}$ defined on the CW-complex *-algebra $\exp(\odot \mathcal{F}_S^{\mathbb{C}})$ and the corresponding state μ defined on the polynomial algebra $\mathbb{C}[\tau]$ we construct now the corresponding GNS representation.

We start with the application of the GNS construction to the state $\mu : \mathbb{C}[\tau] \rightarrow \mathbb{C}$. Consider the corresponding GNS representation $(\mathcal{H}_\mu^0, \langle \cdot, \cdot \rangle_\mu, \rho_\mu, \Omega_\mu)$ of Section 2.4. In particular, the unitary space is

$$\mathcal{H}_\mu^0 = \mathbb{C}[\tau]/J_\mu, \quad (4.28)$$

equipped with the unitary scalar product

$$\langle [P_1], [P_2] \rangle_\mu := \mu(\overline{P_1} P_2). \quad (4.29)$$

Since the state $\tilde{\omega}$ on the algebra $\exp(\odot \mathcal{F}_S^\mathbb{C})$ is the pullback obtained by the epimorphism $\text{Eul} : \exp(\odot \mathcal{F}_S^\mathbb{C}) \rightarrow \mathbb{C}[\tau]$, there is the natural identification

$$(\mathcal{H}_\omega^0, \langle \cdot, \cdot \rangle_\omega) = (\mathcal{H}_\mu^0, \langle \cdot, \cdot \rangle_\mu). \quad (4.30)$$

And the action of the GNS representation ρ_ω becomes:

$$\begin{aligned} \rho_\omega : \exp(\odot \mathcal{F}_S^\mathbb{C}) &\rightarrow \text{End}(\mathcal{H}_\mu^0), \\ \rho_\omega(A)P &:= [\text{Eul}(A) P] \end{aligned} \quad (4.31)$$

using the map (4.25,4.26). Obviously,

$$\Omega_\omega = \Omega_\mu = [1], \quad (4.32)$$

meaning the (equivalence class of) the 0-order polynomial 1.

Proposition 4.14. *$(\mathcal{H}_\mu^0, \langle \cdot, \cdot \rangle_\mu, \rho_\omega, \Omega_\mu)$ defined by (4.28,4.29,4.31,4.32) above is (equivalent to) the GNS representation (2.4) corresponding to the state $\tilde{\omega}$.*

5 $\mathfrak{A}$ defined by the smearing characteristic functions

In this section we combine the definitions and results of the previous Section 4 with the notion of the the polymer *-algebra of Section 2. We consider here the polymer *-algebra $\mathfrak{A}$ defined by the space of the smearing functions $\mathcal{F}$ taken to be the CW-complex vector space $\mathcal{F}_S$,

$$\mathcal{F} = \mathcal{F}_S. \quad (5.1)$$

At the first sight it may be surprising, that in our definition we admit also the smearing functions supported on measure zero lower dimensional simplexes. However, we remember from the previous section, that the simplex vector space $\mathcal{F}_S$ is spanned by the characteristic functions of the open balls in M . Therefore the emergence of the measure zero supports is a result of the linear structure. Still, a state on the corresponding polymer *-algebra $\mathfrak{A}$ might just ignore (in the suitable sense) the quantum momentum variables corresponding to those measure zero sets. We are not making that assumption, though.

We will construct now all the homeomorphism invariant (in the sense of Definition 2.10) states on $\mathfrak{A}$ which satisfy Property (2.10). In the next sections, we will study the corresponding GNS representations, their equivalence classes, and the irreducibility.

The momentum variable space $\Pi_{\mathcal{F}_S}$ (see Definition 2.4) corresponding to that space of the smearing functions will be denoted by

$$\Pi := \Pi_{\mathcal{F}_S}.$$

Since the momentum variables are labeled by the smearing functions, and the smearing functions are labeled in this section by (suitable) subsets in M , we will often use the following notation for the momentum variables assigned to a k -simplex S in M (or more generally, a triangulable subset $S \subset M$),

$$\pi_S := \pi(\mathbf{1}_S), \quad \hat{\pi}_S := \hat{\pi}(\mathbf{1}_S) \quad (5.2)$$

Definition 5.1. *The subalgebra $\mathfrak{P}$ of the polymer $*$ -algebra $\mathfrak{A}$ generated by all the quantum momentum variables $\hat{\pi}(s)$ where s ranges the whole space $\mathcal{F}_S$ of the smearing functions is called the quantum momentum algebra.*

The quantum momentum algebra $\mathfrak{P}$ is naturally isomorphic to the CW-complex algebra $\exp(\odot \mathcal{F}_S^{\mathbb{C}})$,

$$\exp(\odot \mathcal{F}_S^{\mathbb{C}}) \rightarrow \mathfrak{P} \quad (5.3)$$

$$\mathbf{1}_{S_1} \odot \cdots \odot \mathbf{1}_{S_n} \mapsto \hat{\pi}_{S_1} \dots \hat{\pi}_{S_n}, \quad (5.4)$$

where the assignment defined for all the n -tuples of k -simplexes S_i , $i = 1, \dots, n$, $\in \mathbb{N}$, determines the isomorphism.

Now, every homeomorphism invariant state $\omega : \mathfrak{A} \rightarrow \mathbb{C}$ defines by the restriction a homeomorphism invariant state on $\mathfrak{P}$. Therefore, due to Proposition 4.4, for every two balls B_1, B_2 in M ,

$$\omega((\hat{\pi}_{B_1} - \hat{\pi}_{B_2})^2) = 0, \quad (5.5)$$

and since $(\hat{\pi}_B)^* = \hat{\pi}_B$ for every ball B , the equality implies that for every $\hat{a} \in \mathfrak{A}$,

$$\omega(\hat{a}(\hat{\pi}_{B_1} - \hat{\pi}_{B_2})) = 0 = \omega((\hat{\pi}_{B_1} - \hat{\pi}_{B_2})\hat{a}). \quad (5.6)$$

This property leads us to the following observation:

Lemma 5.2. *Every homeomorphism invariant state ω on $\mathfrak{A}$ vanishes on the following elements of $\mathfrak{A}$*

$$\omega(\hat{h}_{k_1, x_1}) = \omega\left(\prod_{j=1}^n \hat{h}_{k_j, x_j}\right) = \omega\left(\prod_{j=1}^n \hat{h}_{k_j, x_j} \prod_{l=1}^m \hat{\pi}(f_l)\right) = 0, \quad (5.7)$$

where all $k_j \neq 0$, the points $x_j \in M$ are pairwise different and $n > 0$.

Proof. Consider

$$\hat{a} = \prod_{j=1}^n \hat{h}_{k_j, x_j} \prod_{l=1}^m \hat{\pi}(f_l) \quad (5.8)$$

such that all $k_j \neq 0$, the points $x_j \in M$ are pairwise different and $n > 0$. It is possible to find two balls B_1 and B_2 in M , such that B_2 does not contain any of the points x_j , $j = 1, \dots, n$, whereas B_1 contains exactly one of the points, say x_1 . Then,

$$[\hat{\pi}_{B_2} - \hat{\pi}_{B_1}, \hat{a}] = k_1 \hat{a}.$$

Acting with ω on the both sides of the above equation and taking advantage of the fact that $\hat{\pi}_{B_2} - \hat{\pi}_{B_1}$ is self-adjoint element of $\mathfrak{A}$ we get

$$0 = \omega((\hat{\pi}_{B_2} - \hat{\pi}_{B_1}) \hat{a}) - \omega(\hat{a} (\hat{\pi}_{B_2} - \hat{\pi}_{B_1})) = k_1 \omega(\hat{a}). \quad (5.9)$$

□

But, the action of every state $\omega : \mathfrak{A} \rightarrow \mathbb{C}$ which satisfies the Property (2.10) is determined by the action on the elements used in Lemma 5.2 and the restriction of ω to the quantum momentum variable subalgebra $\mathfrak{P}$. Therefore, the following is true:

Lemma 5.3. *Every homeomorphism invariant state ω on $\mathfrak{A}$ which satisfies Property (2.10) is determined by its restriction to the quantum momentum subalgebra $\mathfrak{P}$.*

The converse is also true:

Lemma 5.4. *For every state $\tilde{\omega}$ on the quantum momentum subalgebra $\mathfrak{P}$ there exists a unique extension to a homeomorphism invariant state ω of Property 2.10 defined on the polymer *-algebra $\mathfrak{A}$.*

Proof. It is convenient in what follows, to use the quotient algebra $\tilde{\mathfrak{A}}$ (2.41), and the canonical epimorphism $\mathfrak{A} \rightarrow \tilde{\mathfrak{A}}$.

The quantum momentum subalgebra $\mathfrak{P}$ is naturally isomorphic to its image $\tilde{\mathfrak{P}}$ upon the map $\mathfrak{A} \rightarrow \tilde{\mathfrak{A}}$ therefore the subalgebras and their elements will be identified.

Every state $\tilde{\omega}$ on the subalgebra $\mathfrak{P}$, there defines a unique $\mathbb{C}$ -linear extension

$$\tilde{\omega} : \tilde{\mathfrak{A}} \rightarrow \mathbb{C},$$

given by the formulae (5.7) (where all $k_j \neq 0$, the points $x_j \in M$ are pairwise different and $n > 0$) with ω replaced by $\tilde{\omega}$ and the hats replaced by the tildes.

It remains to show that the extension is non-negative. The extension $\tilde{\omega}$ is just the pullback with respect to the following $\mathbb{C}$ -linear map:

$$P : \tilde{\mathfrak{A}} \rightarrow \mathfrak{P} \quad (5.10)$$

$$\begin{aligned} P\left(\prod_{j=1}^n \tilde{\pi}_{U_j}\right) &= \prod_{j=1}^n \hat{\pi}_{U_j} \\ P\left(\prod_{j=1}^n \tilde{h}_{k_j, x_j} \prod_{l=1}^m \tilde{b}\right) &= 0, \end{aligned} \quad (5.11)$$

whenever $k_j \neq 0$, the points $x_j \in M$ are pairwise different and $\tilde{b} \in \mathfrak{P}$. The point is, that the map P is non-negative, in the sense that for every $\tilde{a} \in \tilde{\mathfrak{A}}$ there is $\hat{b}_1, \dots, \hat{b}_n \in \mathfrak{P}$, such that

$$P(\tilde{a}^* \tilde{a}) = \sum_{j=1}^n \hat{b}_j^* \hat{b}_j. \quad (5.12)$$

Indeed, every $\tilde{a} \in \tilde{\mathfrak{A}}$ can be written as a finite sum

$$\tilde{a} = \sum_{j=1}^n \tilde{h}_{\mathbf{k}_j} \tilde{b}_j \quad (5.13)$$

where $\mathbf{k}_j : M \rightarrow \mathbb{R}$, for every $j = 1, \dots, n$ is a finitely supported function (see (2.44)).

Then $P(\tilde{a}^* \tilde{a})$ is given by (5.12).

Therefore

$$\tilde{\omega} = P^* \tilde{\omega} \quad (5.14)$$

is non-negative as well and defines a state on $\mathfrak{A}$ via the pullback $\mathfrak{A} \rightarrow \tilde{\mathfrak{A}}$. $\square$

Combining Lemma 5.3 and Lemma 5.4 with Theorem 4.13 leads to a complete characterization of the homeomorphism invariant states on the polymer *-algebra $\mathfrak{A}$ provided Property 2.10 is satisfied. The states can be labeled by states on the polynomial algebra $\mathbb{C}[\tau]$ (we recall, that the following algebras are naturally identified: $\exp(\odot \mathcal{F}_S^{\mathbb{C}}) = \mathfrak{P} = \tilde{\mathfrak{P}}$ the last one being the image of the previous one in $\tilde{\mathfrak{A}}$):

Theorem 5.5. *Suppose $\mathfrak{A}$ is the polymer *-algebra defined by the space of the smearing functions (5.1). There is a natural bijection between the set of the homeomorphism invariant states on $\mathfrak{A}$ which satisfy Property 2.10 and the set of all the states on the polynomial algebra $\mathbb{C}[\tau]$. The bijection is the pull-back defined by the linear map $\mathfrak{A} \rightarrow \tilde{\mathfrak{A}} \rightarrow \mathbb{C}[\tau]$ where $\mathfrak{A} \rightarrow \tilde{\mathfrak{A}}$ is the canonical projection on the quotient (2.41), whereas the*

map $\tilde{\mathcal{E}} : \tilde{\mathfrak{A}} \rightarrow \mathbb{C}[\tau]$ is defined as follows:

$$\tilde{\mathcal{E}}|_{\mathfrak{P}} = \text{Eul} \quad (5.15)$$

$$\tilde{\mathcal{E}}\tilde{h}_{\mathbf{k}} = 0, \quad (5.16)$$

(where we use the notation of (4.25,4.26,(2.44)) for every non-zero finitely supported function $\mathbf{k}$ on M).

6 Explicit form of GNS representations of $\mathfrak{A}$

We characterize, in this section, the GNS representations (see of the states described by Theorem 5.5. We use the notation and definitions introduced in Section 2.4.

Let ω be a homeomorphism invariant state on $\mathfrak{A}$ whose GNS representation ρ_ω satisfies Property 2.10. Recall, that according to Theorem 5.5, the state ω is the pullback of a state μ on the polynomial algebra $\mathbb{C}[\tau]$. We will also use, that, as explained in Section 2.5, ω is the pullback of a state $\tilde{\omega}$ on the algebra $\tilde{\mathfrak{A}}$ (see (2.41)).

The first step is a characterization of the unitary space $(\mathcal{H}_\omega^0, \langle \cdot, \cdot \rangle_\omega)$ in which the representation is defined; that is, the quotient space $\mathfrak{A}/J_\omega$ endowed with the unitary scalar product $\langle \cdot, \cdot \rangle_\omega$. For this purpose, we will use the unitary spaces: $(\text{Cyl}, (\cdot, \cdot)_{\text{Cyl}})$ (3.11,3.12) and $(\mathbb{C}[\tau]/J_\mu, \langle \cdot, \cdot \rangle_\mu)$ (4.28,4.29). We will also use the $*$ -algebra homomorphism Eul (4.26,4.25) defined also on the quantum momentum algebra

$$\text{Eul} : \mathfrak{P} \rightarrow \mathbb{C}[\tau],$$

via the isomorphism (5.3).

Lemma 6.1. *There is a natural vector space isomorphism*

$$\mathcal{I} : \mathfrak{A}/J_\omega \rightarrow \text{Cyl} \otimes \mathbb{C}[\tau]/J_\mu, \quad (6.1)$$

unitary with respect to the unitary forms $\langle \cdot, \cdot \rangle_\omega$ and, respectively, $(\cdot | \cdot) \otimes \langle \cdot, \cdot \rangle_\mu$, and such that, for every element $\hat{b}$ of the quantum momentum algebra $\mathfrak{P}$ (see Definition 5.1) and every n -tuple of quantum position variables $\hat{h}_{\mathbf{k}_1, x_1}, \dots, \hat{h}_{\mathbf{k}_n, x_n}$, including $\hat{h}_{0, x_i} = \hat{1}$,

$$\mathcal{I} : [\prod_{i=1}^n \hat{h}_{\mathbf{k}_i, x_i} \hat{b}] \mapsto |\sum_{i=1}^n k_i \mathbf{1}_{x_i}\rangle \otimes [\text{Eul}(\hat{b})]. \quad (6.2)$$

Proof. We know that the vector space $\mathfrak{A}/J_\omega$ is isomorphic to $\tilde{\mathfrak{A}}/J_{\tilde{\omega}}$ (see (2.5)). A general element $\tilde{a} \in \tilde{\mathfrak{A}}$ can be written in the form (5.13). Consider the following map,

$$\begin{aligned} \tilde{\mathfrak{A}} &\rightarrow \text{Cyl} \otimes \mathbb{C}[\tau]/J_\mu \\ \sum_{j=1}^n \tilde{h}_{\mathbf{k}_j} \tilde{b}_j &\mapsto \sum_{j=1}^n |\mathbf{k}_j\rangle \otimes [\text{Eul}(\tilde{b}_j)], \end{aligned} \quad (6.3)$$

(as in the previous section, the subalgebra $\mathfrak{P} \subset \mathfrak{A}$ is identified with its image in $\tilde{\mathfrak{A}}$). The kernel of (6.3) is given by the left hand side such that

$$\mathbb{C}[\tau]/J_\mu \ni [\text{Eul}(\tilde{b}_j)] = 0, \quad j = 1, \dots, n. \quad (6.4)$$

On the other hand

$$\tilde{\omega} \left(\left(\sum_{j=1}^n \tilde{h}_{\mathbf{k}_j} \tilde{b}_j \right)^* \sum_{j=1}^n \tilde{h}_{\mathbf{k}_j} \tilde{b}_j \right) = \sum_{j=1}^n \mu((\text{Eul}(\tilde{b}_j))^* \text{Eul}(\tilde{b}_j)). \quad (6.5)$$

Therefore the kernel of (6.3) coincides with the left ideal $J_{\tilde{\omega}}$. Obviously, the map (6.2) is onto. Therefore, it passes to a vector space isomorphism, the isomorphism $\mathcal{I}$.

The calculation (6.5) shows also that $\mathcal{I}$ is unitary, if $\mathfrak{A}/J_\omega$ is endowed with the unitary scalar product $\langle \cdot, \cdot \rangle_\omega$ and the space $\text{Cyl} \otimes \mathbb{C}[\tau]/J_\mu$ with the tensor product $(\cdot|\cdot)_{\text{Cyl}} \otimes \langle \cdot, \cdot \rangle_\mu$. $\square$

Next, we turn to the GNS representation ρ_ω of $\mathfrak{A}$ defined by the state ω . Our task amounts to evaluating the action of the operators $\mathcal{I} \circ \rho_\omega(\hat{h}_{k,x}) \circ \mathcal{I}^{-1}$, and $\mathcal{I} \circ \rho_\omega(\hat{\pi}(s)) \circ \mathcal{I}^{-1}$ in $\text{Cyl} \otimes \mathcal{H}_\mu$. For every element $|\mathbf{k}\rangle \otimes [P]$, we will denote

$$\mathcal{I}^{-1}|\mathbf{k}\rangle \otimes [P] =: [\tilde{h}_{\mathbf{k}} \tilde{b}_P].$$

Given any quantum position variable $\hat{h}_{k,x}$, we have

$$\begin{aligned} \mathcal{I} \circ \rho_\omega(\hat{h}_{k,x}) \circ \mathcal{I}^{-1}|\mathbf{k}\rangle \otimes [P] &= \mathcal{I} \circ [\tilde{h}_{k,x} \tilde{h}_{\mathbf{k}} \tilde{b}_P] = \\ \mathcal{I}[\tilde{h}_{\mathbf{k}+k\mathbf{1}_{\{x\}}} \tilde{b}_P] &= |\mathbf{k} + k\mathbf{1}_{\{x\}}\rangle \otimes [P]. \end{aligned} \quad (6.6)$$

And, given any embedded CW-complex $s \in \mathcal{F}_S$

$$\begin{aligned} \mathcal{I} \circ \rho_\omega(\hat{\pi}(s)) \circ \mathcal{I}^{-1}|\mathbf{k}\rangle \otimes [P] &= \mathcal{I} \circ [\tilde{\pi}(s) \tilde{h}_{\mathbf{k}} \tilde{b}_P] = \\ \mathcal{I}[\tilde{h}_{\mathbf{k}} (\tilde{b}_{\text{Eul}(s)} P + \sum_{x \in M} \mathbf{k}(x) s(x) \tilde{b}_P)] &= |\mathbf{k}\rangle \otimes \left(\sum_{x \in M} \mathbf{k}(x) s(x) [P] + [\text{Eul}(s) P] \right). \end{aligned} \quad (6.7)$$

In conclusion:

Theorem 6.2. *Suppose $\mathfrak{A}$ is the polymer $*$ -algebra defined by the space of the smearing functions (5.1). Suppose ω is a homeomorphism invariant state on $\mathfrak{A}$ such that Property 2.10 is satisfied. Let $\mu : \mathbb{C}[\tau] \rightarrow \mathbb{C}$ be the state corresponding to ω by Theorem 5.5. The GNS representation $(\mathcal{H}_\omega^0, \langle \cdot, \cdot \rangle_\omega, \rho_\omega, \Omega_\omega)$ corresponding to ω (see Section 2.4) is equivalent to $(D, \langle \cdot, \cdot \rangle, \rho, \Omega)$ where (see (3.11, 3.12) and (4.28, 4.29)):*

$$D = \text{Cyl} \otimes \mathcal{H}_\mu^0, \quad (6.8)$$

equipped with the unitary scalar product,

$$\langle \cdot, \cdot \rangle = (\cdot | \cdot)_{\text{Cyl}} \otimes \langle \cdot, \cdot \rangle_\mu, \quad (6.9)$$

and with the distinguished vector,

$$\Omega = |0\rangle \otimes [1], \quad (6.10)$$

and the representation ρ is defined as follows,

$$\rho(\hat{h}_{k,x})|\mathbf{k}\rangle \otimes [P] = |\mathbf{k} + k\mathbf{1}_{\{x\}}\rangle \otimes [P]. \quad (6.11)$$

$$\rho(\hat{\pi}(s))|\mathbf{k}\rangle \otimes [P] = |\mathbf{k}\rangle \otimes \left(\sum_{x \in M} \mathbf{k}(x)s(x)[P] + (-1)^{\dim M} \chi_E(s) \rho_\mu(\tau)[P] \right), \quad (6.12)$$

where GNS representation $(\mathcal{H}_\mu^0, \langle \cdot, \cdot \rangle_\mu, \rho_\mu, \Omega_\mu)$ corresponds to the state $\mu : \mathbb{C}[\tau] \rightarrow \mathbb{C}$.

- Henceforth we will be identifying each GNS representation $(\mathcal{H}_\omega^0, \langle \cdot, \cdot \rangle_\omega, \rho_\omega, \Omega_\omega)$ considered in Theorem 6.2 with the corresponding representation $(D, \langle \cdot, \cdot \rangle, \rho, \Omega)$ defined therein.

7 Self-adjoint extensions

7.1 Momentum-self-adjoint representations

We turn now, to the issue of self-adjoint extension of the GNS representations of the states on the polymer $\ast$ -algebra $\mathfrak{A}$ considered in Section 5 and Section 6. Suppose $\omega : \mathfrak{A} \rightarrow \mathbb{C}$ is a homeomorphism invariant state considered in Theorem 5.5 and determined by a state $\mu : \mathbb{C}[\tau] \rightarrow \mathbb{C}$ on the polynomial algebra. We will study the corresponding representation $(D, \langle \cdot, \cdot \rangle, \rho, \Omega)$ of (6.8 – 6.11) equivalent to the GNS representation corresponding to the state ω . We will also use the Hilbert space $\mathcal{H}$ defined by the completion

$$\mathcal{H} := \bar{D}, \quad (7.1)$$

as well as the GNS representation $(\mathcal{H}_\mu^0, \langle \cdot, \cdot \rangle_\mu, \rho_\mu, \Omega_\mu)$ corresponding to the state $\mu : \mathbb{C}[\tau] \rightarrow \mathbb{C}$ and the Hilbert space

$$\mathcal{H}_\mu = \overline{\mathcal{H}_\mu^0}. \quad (7.2)$$

For every $\hat{a} \in \mathfrak{A}$ the operator $\rho(\hat{a})$ is defined in the domain $D \subset \mathcal{H}$. We know that for every (real valued) smearing function $s \in \mathcal{F}_S$, by the GNS construction, the operator $\rho(\hat{\pi}(s))$ is symmetric. That is, for every pair of vectors $[\hat{a}], [\hat{b}] \in D$

$$\left\langle \rho(\hat{\pi}(s))[\hat{a}], [\hat{b}] \right\rangle = \left\langle [\hat{a}], \rho(\hat{\pi}(s))[\hat{b}] \right\rangle.$$

It is easy to notice, that every operator $\rho(\hat{\pi}(s))$ is essentially self-adjoint if

$$\chi_E(s) = 0. \quad (7.3)$$

Indeed, given $s \in \mathcal{F}_S$ such that (7.3), we have

$$\rho(\hat{\pi}(s))|\mathbf{k}\rangle \otimes [P] = \sum_{x \in M} \mathbf{k}(x)s(x) |\mathbf{k}\rangle \otimes [P],$$

for every finitely supported function $\mathbf{k}$ and every polynomial $P \in \mathbb{C}[\tau]$. The eigenvectors $|\mathbf{k}\rangle \otimes [P]$ span the domain D of the operator, which shows the essential self-adjointness.

Now we will consider the case

$$\chi_E(s) \neq 0.$$

The following theorem shows that the issue of the self-adjointness of the quantum momentum operators $\rho(\hat{\pi}(s))$ comes down to the self-adjointness of the corresponding GNS operators $\rho_\mu(\tau)$ defined in the Hilbert space $\mathcal{H}_\mu$ spanned by polynomials.

Theorem 7.1. *Let $(D, \langle \cdot, \cdot \rangle, \rho, \Omega)$ be any of the representations (6.8-6.11) of the polymer $*$ -algebra $\mathfrak{A}$. The following two conditions are equivalent:*

1. *For every smearing function $s \in \mathcal{F}_S$, the operator $\rho(\hat{\pi}(s))$ is essentially self-adjoint in the Hilbert space $\mathcal{H}$ (7.1);*
2. *the operator $\rho_\mu(\tau)$ is essentially self-adjoint in the Hilbert space $\mathcal{H}_\mu$ (7.2).*

Proof. We will show, that for every $s \in \mathcal{F}_S$, the images of the operators $\rho(\hat{\pi}(s)) \pm i$ are dense in the Hilbert space $\mathcal{H}$, that is, the equalities

$$\overline{(\rho(\hat{\pi}(s)) \pm i)D} = \mathcal{H}, \quad (7.4)$$

are true, if and only if

$$\overline{(\rho_\mu(\tau) \pm i)\mathcal{H}_\mu^0} = \mathcal{H}_\mu. \quad (7.5)$$

The last condition (7.5) is necessary and sufficient for the essential self adjointness of the operator $\rho_\mu(\tau)$.

First step of the proof is to fix an arbitrary vector $|\mathbf{k}\rangle \in \text{Cyl}$, and apply the operator in question to the subspace

$$\{|\mathbf{k}\rangle\} \otimes \mathcal{H}_\mu^0 \subset \text{Cyl} \otimes \mathcal{H}_\mu^0 = D. \quad (7.6)$$

The image is

$$(\rho(\hat{\pi}(s)) \pm i)\{|\mathbf{k}\rangle\} \otimes \mathcal{H}_\mu^0 = \{|\mathbf{k}\rangle\} \otimes ((-1)^M \chi_E(s) \rho_\mu(\tau) + \sum_{x \in M} k(x)s(x) \pm i)\mathcal{H}_\mu^0. \quad (7.7)$$

Focus attention on the second factor of the tensor product on the right hand side. Suppose

$$\chi_E(s) \neq 0. \quad (7.8)$$

The factor is dense in the Hilbert space $\mathcal{H}_\mu$, if and only if the operator $\rho_\mu(\tau)$ is essentially self-adjoint. (On the other hand, if we assume that $\chi_E(s) = 0$, then the second factor is unconditionally dense in $\mathcal{H}_\mu$.)

Now,

$$(\rho(\hat{\pi}(s) \pm i) D = \bigoplus_{|\mathbf{k}\rangle \in \text{Cyl}} \{|\mathbf{k}\rangle\} \otimes ((-1)^M \chi_E(s) \rho_\mu(\tau) + \sum_{x \in M} k(x)s(x) \pm i) \mathcal{H}_\mu^0. \quad (7.9)$$

Clearly, in the case (7.8) the right hand side is dense in $\mathcal{H}$ if and only if the operator $\rho_\mu(\tau)$ is essentially self adjoint. (If $\chi_E(s) = 0$, the right hand side equals D for every ω and μ). $\square$

Definition 7.2. *Every representation $(D, \langle \cdot, \cdot \rangle, \rho, \Omega)$ of the polymer $*$ -algebra defined in (6.8 - 6.11) which satisfies the conditions 1 and 2 will be called momentum self adjoint.*

Theorem 7.1 provides a unique self-adjoint extension for each of the quantum momentum operators, provided certain conditions are satisfied. Talking about representations, however, we need a suitable extension of the common domain of *all* the operators of a given GNS representation. The appropriate framework can be found in [24]. We use some elements of that framework in the next subsection and combine with our very results.

7.2 Unique self-adjointed extension

Definition 7.3. *A hermitian representation of a $*$ -algebra $\mathfrak{A}$ is a triple $(D, \langle \cdot, \cdot \rangle, \rho)$ which consist of*

- *a vector space D equipped with a unitary scalar product $\langle \cdot, \cdot \rangle$,*
- *an algebra homomorphism $\rho : \mathfrak{A} \rightarrow \text{End}(D)$ into the algebra of linear operators defined in D ,*

such that

$$\langle \phi, \rho(a)\psi \rangle = \langle \rho(a^*)\phi, \psi \rangle, \quad \forall \phi, \psi \in D.$$

Given a hermitian representation $(D, \langle \cdot, \cdot \rangle, \rho)$ we also consider the natural Hilbert space completion

$$\mathcal{H} = \bar{D}$$

equipped with the natural extension of the unitary scalar product denoted by the same symbol $\langle \cdot, \cdot \rangle$.

We call a hermitian representation $(D', \langle \cdot, \cdot \rangle', \rho')$ of an algebra $\mathfrak{A}$ an extension of a representation $(D, \langle \cdot, \cdot \rangle, \rho)$ of $\mathfrak{A}$ if: (i) $D \subset D'$, and (ii) $\langle \cdot, \cdot \rangle'$ is an extension of $\langle \cdot, \cdot \rangle$, and (iii)

$$\rho'(a)|_D = \rho(a), \quad \forall a \in \mathfrak{A}. \quad (7.10)$$

For every operator $\rho(a) : D \rightarrow D$ defined by a hermitian representation, the domain D_a^* of the adjoint operator $\rho(a)^*$ is an extension of D ,

$$D \subset D_a^* \subset \mathcal{H}.$$

The operator $\rho(a)$ naturally extends itself to the operator adjoint to $\rho(a^*)$,

$$\rho(a^*)^* : D_a^* \rightarrow \mathcal{H}.$$

Whereas D_a^* may be not preserved by the operator, it can be easily shown that the following intersection is preserved

$$D^* := \bigcap_{b \in \mathfrak{A}} D_b^*, \quad (7.11)$$

$$\rho(a^*)^*(D^*) \subset D^*, \quad \forall a \in \mathfrak{A}. \quad (7.12)$$

Definition 7.4. [24] Given a hermitian representation $(D, \langle \cdot, \cdot \rangle, \rho)$ of a $*$ -algebra $\mathfrak{A}$, the triple $(D^*, \langle \cdot, \cdot \rangle, \rho^*)$ is called the adjoint extension of $(D, \langle \cdot, \cdot \rangle, \rho)$, if

- D^* is defined by (7.11),
- for every $a \in \mathfrak{A}$, $\rho^*(a) : D^* \rightarrow D^*$ is the following linear operator

$$\rho^*(a) := \rho(a^*)^*|_{D^*}. \quad (7.13)$$

It is not hard to check, that the adjoint extension of a hermitian representation of a $*$ -algebra $\mathfrak{A}$ is a representation of $\mathfrak{A}$ in the sense that

$$\rho^*(ab) = \rho^*(a)\rho^*(b).$$

In general though, we do not know if $(D^*, \langle \cdot, \cdot \rangle, \rho^*)$ is hermitian. That property can be ensured by some stronger conditions.

Lemma 7.5. [24] Let $(D, \langle \cdot, \cdot \rangle, \rho)$ be a hermitian representation of a $*$ -algebra $\mathfrak{A}$, and $(D^*, \langle \cdot, \cdot \rangle, \rho^*)$ be its adjoint extension. Let $A \subset \mathfrak{A}$ be the set of elements $a \in \mathfrak{A}$ such that

$$\langle v, \rho^*(a)w \rangle = \langle \rho(a^*)v, w \rangle, \quad \forall v, w \in D^*. \quad (7.14)$$

Suppose A generates the algebra $\mathfrak{A}$. Then $(D^*, \langle \cdot, \cdot \rangle, \rho^*)$ is a hermitian representation of the algebra $\mathfrak{A}$.

We apply now the definitions formulated above and Lemma 7.5 to the representations of the polymer $*$ -algebra. Obviously, each of the representations $(D, \langle \cdot, \cdot \rangle, \rho, \Omega)$ defined by (6.8 - 6.11) of Theorem 6.2 defines just a hermitian representation $(D, \langle \cdot, \cdot \rangle, \rho)$ of the polymer $*$ -algebra (by skipping the cyclic vector Ω). Consider the adjoint extension $(D^*, \langle \cdot, \cdot \rangle, \rho^*)$. In that case the set $A \subset \mathfrak{A}$ introduced in Lemma 7.5 contains every quantum position variable $\hat{h}_{k,x}$ and all those quantum momentum variables $\hat{\pi}(s)$ which satisfy $\chi_E(s) = 0$. Indeed, each operator $\rho(\hat{h}_{k,x})$ is unitary in D , whereas every operator $\rho(\hat{\pi}(s))$ is essentially self adjoint. But this is not sufficient to generate all of $\mathfrak{A}$. Nonetheless, in the case of a momentum self-adjoint representation of $\mathfrak{A}$ (according to Definition 7.2), we also have $\hat{\pi}(s) \in A$ for *every* quantum momentum variable. Then, the set A does generate all the algebra $\mathfrak{A}$. Hence, the adjoint extension is a hermitian representation itself. We can also somewhat simplify the definition of the domain D^* in this case. Indeed, each of the domains $D_{\hat{h}_{k,x}}^*$ is just the full Hilbert space $\bar{D} = \mathcal{H}$. It follows, that in (7.11) the elements $b = \hat{\pi}(s_1) \cdot \dots \cdot \hat{\pi}(s_k)$ are sufficient to achieve the set D^* (note that $D_{\rho(a)}^* \cap D_{\rho(b)}^* \subset D_{\rho(a+b)}^*$ and $D_{\rho(\hat{\pi}(s_1) \cdot \dots \cdot \hat{\pi}(s_k))}^* = D_{\rho(\hat{\pi}(s_1) \cdot \dots \cdot \hat{\pi}(s_k) \rho(\hat{h}_{k,x}))}^*$), and we arrive at the following conclusion:

Corollary 7.6. *Let $(D, \langle \cdot, \cdot \rangle, \rho, \Omega)$ be one of the representations (6.8 - 6.11). Suppose $(D, \langle \cdot, \cdot \rangle, \rho, \Omega)$ is momentum self-adjoint according to Definition 7.2. Then, the adjoint extension $(D^*, \langle \cdot, \cdot \rangle, \rho^*)$ of the hermitian representation $(D, \langle \cdot, \cdot \rangle, \rho)$ of $\mathfrak{A}$ is a hermitian representation. Moreover,*

$$D^* = \bigcap_{s_1, \dots, s_n \in \mathcal{F}_S} D_{\hat{\pi}(s_1) \cdot \dots \cdot \hat{\pi}(s_n)}^*. \quad (7.15)$$

We do not need a complete characterization of the spaces $D_{\hat{\pi}(s_1) \cdot \dots \cdot \hat{\pi}(s_N)}^*$ used in Corollary 7.6. However it is easy to note, that all of them contain a useful common subspace (the one in the middle below):

$$D \subset \text{Cyl} \otimes D_\tau \subset D^*, \quad (7.16)$$

where the subspace $D_\tau \subset \mathcal{H}_\mu$ (see (7.2)) defined in the next paragraph, is the domain of an essentially self-adjoint extension of the operator $\rho_\mu(\tau)$.

Now, the Hilbert space $\mathcal{H}_\mu$ is unitary isomorphic with the space $L^2(\mathbb{R}, d\mu)$, where $d\mu$ is a measure² on $\mathbb{R}$ such that:

- a) $\mu(\tau^n) = \int_{\mathbb{R}} \tau^n d\mu(\tau), \quad n = 0, 1, \dots, n, \dots,$
- b) the subspace of polynomials is dense in $L^2(\mathbb{R}, d\mu)$.

²i.e. regular, Borel measure.

This measure $d\mu$ is uniquely determined by the state μ provided μ satisfies the condition 2 of Theorem 7.1. The unitary isomorphism maps $\tau \in \mathbb{C}[\mathbb{R}]$ into the identity function $\tau : \mathbb{R} \rightarrow \mathbb{R}$. Henceforth, we will be identifying the Hilbert spaces $\mathcal{H}_\mu$ and $L^2(\mathbb{R}, d\mu)$. Via that equivalence, the subspace D_τ is

$$D_\tau = \{f \in L^2(\mathbb{R}, d\mu) : \int_{\mathbb{R}} \tau'^{2n} |f(\tau')|^2 d\mu(\tau') < \infty, \forall n \in \mathbb{N}\}. \quad (7.17)$$

Finally:

Definition 7.7. *A hermitian representation of a $*$ -algebra $\mathfrak{A}$ is called self adjoint if it coincides with its own adjoint extension.*

It can be shown (see [24], Section 8.1) that the adjoint extensions $(D^*, \langle \cdot, \cdot \rangle, \rho^*)$ defined in Corolary 7.6 are all self-adjoint hermitian representations.

7.3 Non-unique self-adjointed extensions

We will generalize here, the unique extension of a momentum self-adjoint representation introduced in the previous subsection to a general case. Consider now an arbitrary state $\mu : \mathbb{C}[\tau] \rightarrow \mathbb{C}$, that is do not assume the property 2 of Theorem 7.1. Still, there exists a measure $d\mu^*$ on $\mathbb{R}$ such that the conditions *a) – b)* hold. The measure defines an essentially self-adjoint extension $\rho_{d\mu^*}^{\text{cl}}(\tau)$ of the operator $\rho_\mu(\tau)$. The domain of the extension is the subspace D_τ (7.17). Identifying again the Hilbert spaces $\mathcal{H}_\mu$ and $L^2(\mathbb{R}, d\mu^*)$, the extension is defined in D_τ (7.17) by,

$$(\rho_{d\mu^*}^{\text{cl}}(\tau)f)(\tau') = \tau' f(\tau'). \quad (7.18)$$

therefore we will denote it just by $\hat{\tau}$,

$$\hat{\tau} := \rho_{d\mu^*}^{\text{cl}}(\tau). \quad (7.19)$$

This extension defines naturally the extension of the GNS representation (6.8-6.11) corresponding to μ , to a hermitian representation $(D_{d\mu^*}, \langle \cdot, \cdot \rangle, \rho_{d\mu^*})$, where $D_{d\mu^*} \subset \mathcal{H}$ such that (we skip the subscript $d\mu^*$ at the scalar product because it coincides with the scalar product in $\mathcal{H}$)

- $D_{d\mu^*} = \text{Cyl} \otimes D_\tau$ equipped with the scalar product of $\text{Cyl} \otimes L^2(\mathbb{R}, d\mu^*)$
- the representation is defined as follows,

$$\rho_{d\mu^*}(\hat{h}_{k,x})|\mathbf{k}\rangle \otimes f = |\mathbf{k} + k\mathbf{1}_{\{x\}}\rangle \otimes f, \quad (7.20)$$

$$\rho_{d\mu^*}(\hat{\pi}(s))|\mathbf{k}\rangle \otimes f = |\mathbf{k}\rangle \otimes \left(\sum_{x \in M} \mathbf{k}(x)s(x)f + (-1)^{\dim M} \chi_E(s)\hat{\tau} \cdot f \right). \quad (7.21)$$

In the same way as it was explained in the previous subsection, it can be shown that the adjoint extension of the hermitian representation $(D_{d\mu^*}, \langle \cdot, \cdot \rangle, \rho_{d\mu^*})$ is a hermitian representation and a self-adjoint representation of $\mathfrak{A}$ in the sense of Definition 7.7.

Finally, in the case of a momentum self-adjoint representation (see Definition 7.2), the extension introduced in the current subsection coincides with the unique extension introduced in Section 7.2.

8 Equivalence of representations

Definition 8.1. *Two representations $(D_1, \langle \cdot, \cdot \rangle_1, \rho_1)$ and $(D_2, \langle \cdot, \cdot \rangle_2, \rho_2)$ are called equivalent if there exists a unitary space isomorphism $I: D_1 \rightarrow D_2$ satisfying:*

$$I^{-1} \circ \rho_2 \circ I = \rho_1. \quad (8.1)$$

I is called an intertwining map between the representations.

We will denote by $\bar{I}$ the unique extension of the map I to the completions $\overline{D_1}$ and $\overline{D_2}$ of the domains.

We will study now, the issue of the equivalence in the context of the momentum self-adjoint representations considered in Section 7.2, as well as in the more general context of the representations defined in Section 7.3. To make our results as general as possible, we consider the (self-)adjoint extensions of the representations. We show, that the equivalence issue comes down to the absolute continuity of the measures on $\mathbb{R}$ labeling the representations.

Theorem 8.2. *Let $(D_1, \langle \cdot, \cdot \rangle_1, \rho_1)$ and $(D_2, \langle \cdot, \cdot \rangle_2, \rho_2)$ be the representations of the algebra $\mathfrak{A}$ defined by: (7.20 - 7.21), and the measures $d\mu^* = d\mu_1, d\mu_2$.*

- *Suppose their adjoint extensions $(D_1^*, \langle \cdot, \cdot \rangle_1, \rho_1^*)$ and $(D_2^*, \langle \cdot, \cdot \rangle_2, \rho_2^*)$ are equivalent. Then:*
 - *The measures $d\mu_1$ and $d\mu_2$ are absolutely continuous with respect to each other;*
 - *For every unitary isomorphism $I: D_1^* \rightarrow D_2^*$ such that*

$$I^{-1} \circ \rho_2^* \circ I = \rho_1^* \quad (8.2)$$

there is $h \in L^2(\mathbb{R}, d\mu_2)$ such that I is of the following form for every $|\mathbf{k}\rangle \in \text{Cyl}$, and every $f \in L^2(\mathbb{R}, d\mu_1)$:

$$|\mathbf{k}\rangle \otimes f \mapsto |\mathbf{k}\rangle \otimes hf. \quad (8.3)$$

Moreover,

$$|h|^2 = \frac{d\mu_1}{d\mu_2}. \quad (8.4)$$

- Conversely, if the measures $d\mu_1$ and $d\mu_2$ are absolutely continuous, the map (8.3) with

$$h := \sqrt{\frac{d\mu_1}{d\mu_2}}$$

intertwines the representations $(D_1, \langle \cdot, \cdot \rangle_1, \rho_1)$ and $(D_2, \langle \cdot, \cdot \rangle, \rho_2)$ (not only the adjoint extensions).

Proof. Suppose the representations $(D_1^*, \langle \cdot, \cdot \rangle_1, \rho_1^*)$ and $(D_2^*, \langle \cdot, \cdot \rangle_2, \rho_2^*)$ are equivalent and $\bar{I}: \bar{D}_1 \rightarrow \bar{D}_2$ is an intertwining map. The sketch of the proof of (8.3) is simple. Every intertwining map I is determined by its action on the cyclic vector $|\mathbf{0}\rangle \otimes 1$. But, as we argue below, each intertwining map I has to satisfy

$$I : |\mathbf{0}\rangle \otimes 1 \mapsto |\mathbf{0}\rangle \otimes h, \quad (8.5)$$

where $h \in L^2(\mathbb{R}, d\mu_2)$. Next, we find that the map I determined by (8.5) is exactly the one defined in (8.3). Below we go to the details. The only technical subtlety we have to be careful about is that we are dealing with the adjoint extension of the representation ρ_2 .

The vector $|\mathbf{0}\rangle \otimes 1$ is annihilated by the following sub-class of the momentum operators:

$$\rho_1(\hat{\pi}(s))|\mathbf{0}\rangle \otimes 1 = 0, \quad \text{whenever } \chi_E(s) = 0. \quad (8.6)$$

This property has to be preserved by every intertwiner, therefore

$$\rho_2^*(\hat{\pi}(s))I(|\mathbf{0}\rangle \otimes 1) = 0, \quad \text{whenever } \chi_E(s) = 0. \quad (8.7)$$

But every vector $\psi \in D_2^*$ such that

$$\rho_2^*(\hat{\pi}(s))\psi = 0, \quad \forall s \in \mathcal{F}_S : \chi_E(s) = 0, \quad (8.8)$$

is necessarily of the form $|\mathbf{0}\rangle \otimes h$ (that statement is more obvious in the case of $\psi \in D_2$ and ρ_2 , and it generalizes to the adjoint extensions), hence:

$$I(|\mathbf{0}\rangle \otimes 1) = |\mathbf{0}\rangle \otimes h, \quad h \in L^2(\mathbb{R}, d\mu_2). \quad (8.9)$$

Next we determine the action of I on the vectors $|\mathbf{0}\rangle \otimes \tau^n$, $n = 1, \dots$. We have

$$I(|\mathbf{0}\rangle \otimes \tau^n) = \rho_2^*(\hat{\pi}(s_1)) \cdots \rho_2^*(\hat{\pi}(s_n))|\mathbf{0}\rangle \otimes h, \quad (8.10)$$

with any $s_1, \dots, s_n \in \mathcal{F}_S$ such that $\chi_E(s_i) = 1$, $i = 1, \dots, n$. Again, we observe that the right hand side of (8.10), if denoted by ψ , satisfies (8.8). Hence, for every $s \in \mathcal{F}_S$,

the operator $\rho_2^*(\hat{\pi}(s))$ preserves the subspace $|\mathbf{0}\rangle \otimes L^2(\mathbb{R}, d\mu_2) \cap D_2^*$. Fix s such that $\chi_E(s) = 1$ and consider the restriction

$$\rho_2^*(\hat{\pi}(s)) : |\mathbf{0}\rangle \otimes L^2(\mathbb{R}, d\mu_2) \cap D_2^* \rightarrow |\mathbf{0}\rangle \otimes L^2(\mathbb{R}, d\mu_2) \cap D_2^*. \quad (8.11)$$

It follows from (7.21) and (7.19) that the operator (8.11) coincides with the operator $\hat{\tau}$ (as expected). Therefore, we have derived

$$I(|\mathbf{0}\rangle \otimes \tau^n) = |\mathbf{0}\rangle \otimes \hat{\tau}^n \cdot h = |\mathbf{0}\rangle \otimes h\tau^n. \quad (8.12)$$

Finally, the action of the intertwiner I on elements $|\mathbf{k}\rangle \otimes \tau^n$ is determined in the obvious way:

$$I(|\mathbf{k}\rangle \otimes \tau^n) = I(\rho_1(\prod_{x \in M} \hat{h}_{\mathbf{k}(x),x}))|\mathbf{0}\rangle \otimes \tau^n = \rho_2^*(\prod_{x \in M} \hat{h}_{\mathbf{k}(x),x})|\mathbf{0}\rangle \otimes h\tau^n = |\mathbf{k}\rangle \otimes h\tau^n.$$

By the continuity the map extends to (8.3).

The absolute continuity of the measures $d\mu_1$ and $d\mu_2$ is obvious as well as the Nikodym-Radon derivative (8.4).

Conversely, given two measures $d\mu_1$ and $d\mu_2$ on $\mathbb{R}$ which are absolutely continues, one can check by inspection that the map (8.3) is unitary, maps $D_1 \rightarrow D_2$ and intertwines the representations ρ_1 and ρ_2 . $\square$

9 Invariant subspaces

In this section we solve the issue of the irreducibility of the representations of the polymer *-algebra $\mathfrak{A}$ introduced in (6.8,6.9,6.11) and in Sections 7.2, 7.3.

Consider a representation $(D, \langle \cdot, \cdot \rangle, \rho)$ (6.8,6.9,6.11) of the algebra $\mathfrak{A}$. As before, we will also use the corresponding Hilbert space $\mathcal{H}$ (7.1). It is defined by the completed tensor product (we denote it by $\overline{\otimes}$) of the following spaces

$$\overline{\text{Cyl}}, \quad \mathcal{H}_\mu = \overline{\mathbb{C}[\tau]/J_\mu}, \quad (9.1)$$

the completions with respect to the unitary scalar products $(\cdot|\cdot)_{\text{Cyl}}$, and, respectively $\langle \cdot, \cdot \rangle_\mu$, that is

$$\mathcal{H} = \overline{\text{Cyl}} \overline{\otimes} \mathcal{H}_\mu. \quad (9.2)$$

We present a quite strong, general result which shows that the issue boils down to the irreducibility of the representations of the polynomial algebra $\mathbb{C}[\tau]$ corresponding to the state $\mu : \mathbb{C}[\tau] \rightarrow \mathbb{C}$.

Recall, that each quantum position operator $\rho(\hat{h}_{k,x}) : \mathcal{H} \rightarrow \mathcal{H}$ is an unitary operator after extension to the whole space. Another class of unitary operators can be defined

by using the quantum momentum operators $\rho(\hat{\pi}(s))$ corresponding to those complex-smearing-functions $s \in \mathcal{F}_S$ which satisfy

$$\chi_E(s) = 0. \quad (9.3)$$

Each of those operators with the domain D (6.8) is essentially-self adjoint. Therefore, the operator $\exp(i\rho(\hat{\pi}(s)))$ can be defined uniquely as an unitary operator in $\mathcal{H}$. We will assume below that the unitary operators mentioned above preserve a common subspace in $\mathcal{H}$, and we will come to quite strong conclusion.

Theorem 9.1. *Let $(D, \langle \cdot, \cdot \rangle, \rho)$ be a representation of the polymer $*$ -algebra defined by (6.8, 6.9, 6.11), corresponding to a state $\omega : \mathfrak{A} \rightarrow \mathbb{C}$ determined by a state $\mu : \mathbb{C} \rightarrow \mathbb{C}$. Suppose $\tilde{\mathcal{H}}$ is a Hilbert subspace of the Hilbert space $\mathcal{H}$ (9.2) such that*

$$\rho(\hat{h}_{k,x})(\tilde{\mathcal{H}}) \subset \tilde{\mathcal{H}} \quad (9.4)$$

$$\exp(i\rho(\hat{\pi}(s)))(\tilde{\mathcal{H}}) \subset \tilde{\mathcal{H}}, \quad (9.5)$$

for every quantum position variable $\hat{h}_{k,x}$ and every quantum momentum $\hat{\pi}(s)$ such that (7.3). Then, there is a Hilbert subspace $\tilde{\mathcal{H}}_\mu$ of $\mathcal{H}_\mu$ such that

$$\tilde{\mathcal{H}} = \overline{\text{Cyl}} \otimes \tilde{\mathcal{H}}_\mu. \quad (9.6)$$

Proof. Given $s \in \mathcal{F}_S$, every eigenvalue of the operator $\exp(i\rho_\omega(\hat{\pi}(s)))$ can be written as

$$\lambda_{\mathbf{k},s} = e^{i\sum_{x \in M} \mathbf{k}(x)s(x)}, \quad (9.7)$$

where $\mathbf{k}$ is a finitely supported function on M , and for every $\mathbf{k}$ the number $\lambda_{\mathbf{k},s}$ is an eigenvalue of the operator $\exp(i\rho_\omega(\hat{\pi}(s)))$. To each of the eigenvalues $\lambda_{\mathbf{k},s}$, there is assigned the corresponding subspace $\mathcal{H}_{\lambda_{\mathbf{k},s}}$ of the eigenvectors. We have,

$$|\mathbf{k}'\rangle \otimes \mathcal{H}_\mu \subset \mathcal{H}_{\lambda_{\mathbf{k},s}}, \quad (9.8)$$

for every finitely supported function $\mathbf{k}'$ such that

$$\sum_{x \in M} \mathbf{k}'(x)s(x) = \sum_{x \in M} \mathbf{k}(x)s(x).$$

However, if we fix a finitely supported function $\mathbf{k}$ and consider the common part of all the sets $\mathcal{H}_{\lambda_{\mathbf{k},s}}$, then it is easy to show that

$$\bigcap_{s \in \mathcal{F}_S : \chi_E(s)=0} \mathcal{H}_{\lambda_{\mathbf{k},s}} = |\mathbf{k}\rangle \otimes \mathcal{H}_\mu. \quad (9.9)$$

Let us turn now, to the preserved subspace $\tilde{\mathcal{H}}$. Denote by

$$P_{\tilde{\mathcal{H}}} : \mathcal{H} \rightarrow \tilde{\mathcal{H}}$$

the orthogonal projection onto $\tilde{\mathcal{H}}$. For every fixed $s \in \mathcal{F}_S$ such that (7.3), and a finitely supported function $\mathbf{k}$, due to the assumption that $\tilde{\mathcal{H}}$ is preserved by the unitary operator $\exp(i\rho_\omega(\hat{\pi}(s)))$ and by the general properties of the unitary operators, it follows that

$$P_{\tilde{\mathcal{H}}}(\mathcal{H}_{\lambda_{\mathbf{k},s}}) \subset \mathcal{H}_{\lambda_{\mathbf{k},s}}. \quad (9.10)$$

Therefore

$$P_{\tilde{\mathcal{H}}}(|\mathbf{k}\rangle \otimes \mathcal{H}_\mu) = \bigcap_s P_{\tilde{\mathcal{H}}}(\mathcal{H}_{\lambda_{\mathbf{k},s}}) \subset |\mathbf{k}\rangle \otimes \mathcal{H}_\mu, \quad (9.11)$$

where the range of s on the right hand side is the same as in (9.9). Hence, there is a Hilbert subspace $\tilde{\mathcal{H}}_{\mathbf{k},\mu} \subset \mathcal{H}_\mu$, such that

$$P_{\tilde{\mathcal{H}}}(|\mathbf{k}\rangle \otimes \mathcal{H}_\mu) = |\mathbf{k}\rangle \otimes \tilde{\mathcal{H}}_{\mathbf{k},\mu}. \quad (9.12)$$

The relevance of this space consists in the following orthogonal decomposition of the invariant space $\tilde{\mathcal{H}}_\omega$,

$$\tilde{\mathcal{H}} = P_{\tilde{\mathcal{H}}}(\overline{\bigoplus_{\mathbf{k}} |\mathbf{k}\rangle \otimes \tilde{\mathcal{H}}_\mu}) = \overline{\bigoplus_{\mathbf{k}} |\mathbf{k}\rangle \otimes \tilde{\mathcal{H}}_{\mathbf{k},\mu}}, \quad (9.13)$$

where $\mathbf{k} : M \rightarrow \mathbb{R}$ ranges the set of all the finitely supported functions.

Next we will show, that in fact the subspaces $\tilde{\mathcal{H}}_{\mathbf{k},\mu}$ are necessarily all the same, independent of $\mathbf{k}$. To see that, we introduce for every finitely supported function $\mathbf{k} : M \rightarrow \mathbb{R}$ the following unitary operator,

$$U_{\mathbf{k}} = \prod_{x \in M} \rho(\hat{h}_{\mathbf{k}(x),x}). \quad (9.14)$$

By the definition of the operators $U_{\mathbf{k}}$, and $\rho(\hat{h}_{k,x})$,

$$U_{\mathbf{k}}(|\mathbf{k}'\rangle \otimes \mathcal{H}'_\mu) = |\mathbf{k} + \mathbf{k}'\rangle \otimes \mathcal{H}'_\mu, \quad (9.15)$$

for every two finitely supported functions $\mathbf{k}$ and $\mathbf{k}'$ defined on M , and every Hilbert subspace $\mathcal{H}'_\mu \subset \mathcal{H}_\mu$. On the other hand, the invariant space $\tilde{\mathcal{H}}$ is in particular both $U_{\mathbf{k}}$ and $(U_{\mathbf{k}})^{-1} = U_{-\mathbf{k}}$ invariant, hence

$$U_{\mathbf{k}}(\tilde{\mathcal{H}}) = \tilde{\mathcal{H}}. \quad (9.16)$$

In conclusion,

$$|\mathbf{k}\rangle \otimes \tilde{\mathcal{H}}_{\mathbf{k},\mu} = U_{\mathbf{k}}(|\mathbf{0}\rangle \otimes \tilde{\mathcal{H}}_{\mathbf{k},\mu}) = \quad (9.17)$$

$$U_{\mathbf{k}}(|\mathbf{0}\rangle \otimes \mathcal{H}_{\mu} \cap \tilde{\mathcal{H}}) = U_{\mathbf{k}}(|\mathbf{0}\rangle \otimes \tilde{\mathcal{H}}_{\mathbf{0},\mu}) = \quad (9.18)$$

$$= |\mathbf{k}\rangle \otimes \tilde{\mathcal{H}}_{\mathbf{0},\mu}. \quad (9.19)$$

Therefore, the subspace pointed out in the conclusion of this theorem is

$$\tilde{\mathcal{H}}_{\mu} = \tilde{\mathcal{H}}_{\mathbf{0},\mathbf{k}}.$$

□

To complete the conclusion of Theorem 9.1, consider a hermitian representation $(D, \langle \cdot, \cdot \rangle, \rho)$ of $\mathfrak{A}$. Suppose it is either defined by a momentum self-adjoint GNS representation and by formulae (6.8,6.9,6.11), or it is one of the representations (7.20,7.21). In any case, each of the operators $\rho((\hat{\pi}(s)))$, $s \in \mathcal{F}_{\mathcal{S}}$, is essentially self adjoint, hence it defines a unitary operator $\exp(i\rho(\hat{\pi}(s)))$. Suppose the subspace

$$\tilde{\mathcal{H}} = \overline{\text{Cyl}} \otimes \tilde{\mathcal{H}}_{\mu}$$

of Theorem 9.1 is preserved by each of the operators $\exp(i\hat{\pi}(s))$. Then the subspace

$$\tilde{\mathcal{H}}_{\mu} \subset L^2(\mathbb{R}, d\mu^*)$$

is necessarily preserved by the operator $\exp(i\lambda\hat{\tau})$, $\lambda \in \mathbb{R}$ (see 7.19). On the other hand, generically, it is not hard to find such a subspace $\tilde{\mathcal{H}}_{\mu}$. To see an example, suppose there is a $d\mu^*$ -measurable, proper subset $V \subset \mathbb{R}$. Define a Hilbert subspace $\tilde{\mathcal{H}}_{\mu} \subset L^2(\mathbb{R}, d\mu^*)$ to be spanned by (classes of) all the square integrable functions of supports contained in V . The Hilbert space $\tilde{\mathcal{H}} = \overline{\text{Cyl}} \otimes \tilde{\mathcal{H}}_{\mu}$ has the non-trivial (dense in $\tilde{\mathcal{H}}$) intersection with the domain D^* of the adjoint extension

$$\tilde{D} = D^* \cap \tilde{\mathcal{H}}. \quad (9.20)$$

The subspace $\tilde{D}$ is invariant with respect to the representation ρ^* . Hence, the restriction of ρ^* to $\tilde{D}$ is a new hermitian representation of the algebra $\mathfrak{A}$.

That representation is equivalent to the adjoint extension $(D_{d\mu'}^*, \langle \cdot, \cdot \rangle', \rho_{d\mu'}^*)$ of the representation $(D_{\mu'}, \langle \cdot, \cdot \rangle_{\mu'}, \rho_{\mu'})$ defined by (7.20,7.21) and a measure

$$d\tilde{\mu}' = \mathbf{1}_V d\mu^*.$$

The only case in which there is no invariant subspace $\tilde{\mathcal{H}}$ in Theorem 9.1, is the state $\mu : \mathbb{C} \rightarrow \mathbb{C}$ defined by a measure $d\mu = \delta_{r_0}$ supported at a single point $r_0 \in \mathbb{R}$.

10 Discussion

10.1 Summary of the results

The polymer $*$ -algebra $\mathfrak{A}$ (Definition 2.7) contains the momentum subalgebra generated by the smeared momentum variables (Definition 2.3). The study of diffeomorphism invariant states on the momentum subalgebra is crucial for characterization of possible diffeomorphism invariant states on $\mathfrak{A}$.

In the $C^{(n)}$, compactly supported smearing functions case (Sec. 3), there is no non-trivial $C^{(n)}$ -diffeomorphism invariant state on the momentum subalgebra. This is the meaning of the identity (3.3) derived from the invariance. This fact and the assumption Property 2.10 determine the unique state on $\mathfrak{A}$ (3.2). The corresponding representation (3.13, 3.14) of $\mathfrak{A}$ is the one used in LQG in the quantization of the scalar field. The uniqueness was not a surprise, and the argument used in the proof was similar to that of [23].

The remaining part of the work was focused on the case of the momentum variables $\pi(s)$ defined by smearing against the characteristic functions of regions in M . The result is the construction of new states. Now, the momentum subalgebra -identified with the CW-complex algebra $\exp(\odot \mathcal{F}_S^{\mathbb{C}})$ in this case (Sec. 4.2)- does admit non-trivial, diffeomorphism invariant states. In order to construct explicit examples, we assumed a greater symmetry, namely the homeomorphism invariance (see Sec. 2.6). We succeeded in deriving all the states. The complete class is characterized in Theorem 4.13.

Again, each of the states defined on the momentum subalgebra determines a state on the polymer $*$ -algebra $\mathfrak{A}$ upon the assumption Property 2.10. This leads to the derivation of all the homeomorphism invariant states defined on the polymer $*$ -algebra $\mathfrak{A}$ which satisfy Property 2.10 (Theorem 5.5). The states are labeled by all the states $\mu : \mathbb{C}[\tau] \rightarrow \mathbb{C}$ in a 1-1 manner, where $\mathbb{C}[\tau]$ is the $*$ -algebra of the polynomial functions on $\mathbb{R}$, and $\tau = \text{id} : \mathbb{R} \rightarrow \mathbb{R}$.

The resulting GNS representations of $\mathfrak{A}$ corresponding to the derived states, are provided in explicit form in Theorem 6.2. The properties of the representations are analyzed in Sec. 7 - 9.

Given any of the representations ρ of $\mathfrak{A}$, the momentum operators $\rho(\hat{\pi}(s))$ may or may not be essentially self-adjoint. The necessary and sufficient conditions for that momentum self-adjointness are formulated in terms of the labeling state $\mu : \mathbb{C}[\tau] \rightarrow \mathbb{C}$ in Theorem 7.1.

Whereas the self-adjoint extension of each of the essentially self-adjoint momentum operators is quite well understood, the corresponding extension of the entire representation ρ requires the special care. The suitable framework was derived by Schmudgen [24]. We show that for each of our momentum self-adjoint representations ρ of $\mathfrak{A}$, the adjoint extension ρ^* (Definition 7.4) is a self-adjoint (Definition 7.7) hermitian representation

of $\mathfrak{A}$. The remaining representations we found are also extended to homeomorphism invariant, self-adjoint according to [24], hermitian representations of $\mathfrak{A}$. Each resulting self-adjoint representation $\rho_{d\mu}^*$ of $\mathfrak{A}$ is determined by a measure $d\mu$ on $\mathbb{R}$, provided the algebra of the polynomials is dense in $L^2(\mathbb{R}, d\mu)$. This is the outcome of Sec. 7.

Two self-adjoint representations $\rho_{d\mu_1}^*$ and $\rho_{d\mu_2}^*$ constructed in Sec. 7 are equivalent if and only if the measures $d\mu_1$ and $d\mu_2$ are absolutely continues with respect to each other (Theorem 8.2). The intertwining Hilbert space isomorphism is found as well.

The issue of the reducibility of the representations considered above is solved in Sec. 9. A general characterization of invariant subspaces is given (Theorem 9.1). The final conclusion is:

Corollary 10.1. *Suppose a representation $(\rho, D, \langle \cdot, \cdot \rangle)$ of the polymer $*$ -algebra $\mathfrak{A}$ is either*

- *the representation defined by (6.8, 6.9, 6.11) and a state $\mu : \mathbb{C}[\tau] \rightarrow \mathbb{C}$, and assume it is momentum self-adjoint;*
- *the representations (7.20, 7.21) defined by a measure $d\mu^*$ on $\mathbb{R}$.*

Then, there is no proper subspace $\tilde{\mathcal{H}} \subset \bar{D}$ preserved by the unitary extensions of all the operators $\exp(i\rho(\hat{\pi}(s)))$ and $\rho(\hat{h}_{k,x})$, unless the state μ (the measure $d\mu^$) is the delta measure δ_{τ_0} supported at $\tau_0 \in \mathbb{R}$. The definition of the representation reads:*

$$\begin{aligned} (D, \langle \cdot, \cdot \rangle) &= (\text{Cyl}, (\cdot|_{\text{Cyl}})_{\text{Cyl}}) \\ \rho(\hat{h}_{k,x})|\mathbf{k}\rangle &= |\mathbf{k} + k\mathbf{1}_{\{x\}}\rangle, \\ \rho(\hat{\pi}(s))|\mathbf{k}\rangle &= \left(\sum_{x \in M} \mathbf{k}(x)s(x) + \tau_0(-1)^{\dim M} \chi_E(s) \right) |\mathbf{k}\rangle. \end{aligned} \quad (10.1)$$

If $\tau'_0 \neq \tau_0$, then the representation ρ' corresponding to τ'_0 is inequivalent ρ .

10.2 Relevance for the holonomy-flux algebra

The polymer $*$ -algebra considered in this work is used in Loop Quantum Gravity for the quantum scalar field coupled with the quantum geometry. The quantum geometry itself is described in terms of the holonomy-flux $*$ -algebra [23]. The holonomy-flux $*$ -algebra has a similar structure to the polymer $*$ -algebra, and one could say that the latter is a simplified version of the former one. In particular, the quantum flux variable $\hat{P}(f)$ -the counter part of the quantum momentum operators- is defined by a smearing function $f : M \rightarrow \mathfrak{su}(2)$ of the support contained in an oriented 2-simplex s in a 3-dimensional piecewise-analytic manifold M , and taking values in the Lie algebra $\mathfrak{su}(2)$. In particular, f is often assumed to be of the form

$$f = \mathbf{1}_s \xi, \quad \xi \in \mathfrak{su}(2). \quad (10.2)$$

The corresponding momentum can be denoted by $\hat{P}_{s,\xi}$. Naturally, the question arises, if the class of the topological states defined on the simplex algebra in Section 4 of the current paper can be used to define a class of new diffeomorphism invariant states on the holonomy-flux algebra. Indeed, by exactly the same argument as in Section 4 we prove, that for every topologically invariant state ω

$$\omega((\hat{P}_{s,\xi} - \hat{P}_{s',\xi})^*(\hat{P}_{s,\xi} - \hat{P}_{s',\xi})) = 0, \quad (10.3)$$

for two arbitrary piecewise-analytic 2-simplexes s and s' in M . However, in this case a diffeomorphism which flips the orientation of s into the opposite one is, on the one hand, a symmetry of ω but on the other hand maps

$$\hat{P}(s, \xi) \mapsto -\hat{P}_{s,\xi}. \quad (10.4)$$

It follows from (10.3,10.4) that

$$\omega(\hat{P}_{s,\xi}^* \hat{P}_{s,\xi}) = 0.$$

Eventually, the only state is the one used in LQG. However, if we relaxed the topological invariance assumption, in favor of proper diffeomorphism invariance, perhaps new states could be found on the holonomy-flux algebra defined by the characteristic functions. On the other hand we have the uniqueness result [23] valid in the case of the compactly supported, $C^{(n)}$ -smearing functions f (analogous to Section 3).

A Piecewise-analytic simplexes and manifolds

In the main part of this work we are using piecewise-analytic simplexes embedded in a manifold M endowed with an (appropriately defined) piecewise-analytic structure. Those notions are defined below. The property of the piecewise-analytic simplexes crucial in this paper is Property A.7 below.

Definition A.1. [26] *A subset $X \subset \mathbb{R}^N$ is called semi-analytic if it has the following property: for every $x \in \bar{X}$ (the completion of X) there is a neighborhood $U \ni x$ in $\mathbb{R}^N$, and analytic functions $f_{ij} : U \rightarrow \mathbb{R}$, $i = 1, \dots, n$, $j = 1, \dots, n_i$ such that*

$$\begin{aligned} U \cap X &= \bigcup_{i=1}^n X_i \\ X_i &= \{x \in U : f_{ij}(x) \leq 0, j = 1, \dots, n_i\}. \end{aligned} \quad (A.1)$$

Obviously, every analytic diffeomorphism maps semi-analytic sets into semi-analytic sets. However, the class of the diffeomorphisms of that property is larger. Therefore, we define:

Definition A.2. A $C^{(n)}$ -diffeomorphism $\phi : U \rightarrow U'$, where U , and U' are open subsets in $\mathbb{R}^N$, $0 \leq n < \infty$, is called *piecewise-analytic* if for every semi-analytic subset X of $\mathbb{R}^N$, the subsets $\phi(X \cap U)$ and $\phi^{-1}(X \cap U')$ are semi-analytic.

A sub-family of piecewise-analytic diffeomorphisms was introduced in [23] and called semi-analytic diffeomorphisms. Many examples were constructed. In particular, it was shown, that for every point $x \in \mathbb{R}^N$, and every neighborhood U of x , there is a semi-analytic diffeomorphism which moves x but is the identity map on $\mathbb{R}^N \setminus U$. In this sense the structure we are fixing in $\mathbb{R}^N$ to consider the semi-analytic sets admits local degrees of freedom.

Given a manifold M , a structure compatible with the semi-analytic sets can be defined as follows. Let M be a differentiable (or topological) manifold of the differentiability class $C^{(n)}$, $0 < n < \infty$ (respectively, $n = 0$). Suppose the maximal atlas admits a sub-atlas $((U_I, \chi_I))_{I \in \mathcal{I}}$ labeled by some set $\mathcal{I}$, such that for every two chard χ_I and χ_J , the map

$$\chi_J \circ \chi_I^{-1} : \chi_I(U_I \cap U_J) \rightarrow \chi_J(U_I \cap U_J)$$

is a piecewise-analytic $C^{(n)}$ -diffeomorphism. We call the family $((\chi_I, U_I))_{I \in \mathcal{I}}$ a piecewise analytic atlas on M and extend to a maximal piecewise analytic atlas. A chard (χ, U) belonging to the maximal piecewise-analytic atlas is called a piecewise-analytic chard.

Definition A.3. A manifold endowed with a piecewise analytic atlas is called *piecewise analytic*.

Obviously, every analytic manifold is piecewise analytic. A piecewise analytic diffeomorphism $M \rightarrow M$ of a piecewise analytic manifold M is a diffeomorphism such that itself and its inverse preserve the maximal piecewise analytic atlas.

Now we turn to semi-analytic simplexes. As before, we begin with $\mathbb{R}^N$:

Definition A.4. A 0-simplex in $\mathbb{R}^N$ is a point. If $k > 0$, a piecewise-analytic k -simplex in $\mathbb{R}^N$ is a semi-analytic set S such that there is a homeomorphism $\phi : \mathbb{R}^N \rightarrow \mathbb{R}^N$ which maps S and the completion $\bar{S}$ onto the following sets

$$\phi(S) = \{(x^1, \dots, x^N) \in \mathbb{R}^N : \sum_{i=1}^k x^i < 1, \text{ and } x^{k+1}, \dots, x^N = 0\} \quad (\text{A.2})$$

$$\phi(\bar{S}) = \{(x^1, \dots, x^N) \in \mathbb{R}^N : \sum_{i=1}^k x^i \leq 1, \text{ and } x^{k+1}, \dots, x^N = 0\}. \quad (\text{A.3})$$

And next we proceed with a manifold:

Definition A.5. A k -simplex in a piecewise-analytic manifold is a subset $S \subset M$ such that there is a piecewise-analytic chard (χ, U) which maps S onto a semi-analytic k -simplex in $\mathbb{R}^N$.

The simplexes are used for partitions called triangulations:

Definition A.6. *A piecewise-analytic triangulation of subsets $X_1, \dots, X_n \subset M$, where M is a piecewise-analytic manifold, is a family of pairwise disjoint subsets $S_1, \dots, S_m \subset M$, such that S_i is a k_i -simplex in M for $i = 1, \dots, n$, and*

$$X_i = \bigcup_{k=1}^{n_i} S_{j_{ik}}, \quad i = 1, \dots, n, \quad 1 \leq j_{ik} \leq m. \quad (\text{A.4})$$

The property of the semi-analytic sets and piecewise-analytic triangulations crucial for our work is [26] (see Łojasiewicz, page 463, Theorem 1,2):

Property A.7. *Every finite family of piecewise-analytic simplexes in a piecewise analytic manifold M admits a piecewise-analytic triangulation.*

Let us remark on the status of the framework we are using. The theory of the semi-analytic sets is well established in the mathematical literature [26]. We apply the deep results of that theory. There seems to be no unique generalization of the notion of semi-analyticity to a category of ‘semi-analytic’ manifolds, though. (Probably each generalization has its drawbacks and limitations.) In the current paper we introduce the weakest from the point of view of our aims definition of a manifold structure compatible with the definition of the semi-analytic sets. It relies on a somewhat implicit (but not empty) definition of the piecewise-analytic diffeomorphisms in $\mathbb{R}^N$. In [23] we introduced a family of explicitly defined ‘semi-analytic’ diffeomorphisms, and used them to define ‘semi-analytic’ manifolds. They form a subclass in the class of the piecewise-analytic manifolds defined above. Finally, our definitions should be also compared with similar ideas of [27].

Acknowledgments We have benefited from discussions with Abhay Ashtekar, John Baez, Marcin Bobieński, Witold Marciszewski, Tadeusz Mostowski, Hanno Sahlmann, Lee Smolin, Thomas Thiemann and Andrzej Trautman. The work was partially supported by the Polish Ministerstwo Nauki i Informatyzacji grant 1 P03A 015 29.

References

- [1] Smolin L 2005 The case for background independence [hep-th/0507235]
- [2] Rovelli C and Smolin L 1988 Knot theory and quantum gravity *Phys. Rev. Lett.* **61** 1155-1158
- [3] Rovelli C and Smolin L 1990 Loop representation for quantum general relativity *Nucl. Phys.* **B331** 80–152

- [4] Ashtekar A 1991 *Lectures on non-perturbative canonical gravity* Notes prepared in collaboration with R. S. Tate (World Scientific, Singapore)
- [5] Ashtekar A and Lewandowski J 1994 Representation theory of analytic holonomy algebras, in *Knots and Quantum Gravity* ed Baez J C (Oxford U. Press, Oxford)
- [6] Baez J C 1994 Generalized measures in gauge theory *Lett. Math. Phys.* **31** 213–223
- [7] Lewandowski J 1994 Topological Measure and Graph-Differential Geometry on the Quotient Space of Connections *Int. J. Mod. Phys.* **D3** 207–210
- [8] Lewandowski J. 1997 Volume and Quantizations *Class. Quant. Grav.* **14** 71–76
- [9] Ashtekar A and Lewandowski J 1997 Quantum theory of geometry I: Area operators *Class. Quant. Grav.* **14** A55–A81
- [10] Ashtekar A and Lewandowski J 1997 Quantum theory of geometry II: Volume Operators *Adv. Theo. Math. Phys.* **1** 388–429
- [11] De Pietri R 1997 Spin Networks and Recoupling in Loop Quantum Gravity *Nucl. Phys. Suppl* **57** 251–254
- [12] De Pietri R 1997 On the relation between the connection and the loop representation of quantum gravity *Class. Quant. Grav.* **14** 53–70
- [13] Thiemann T 1998 Closed formula for the matrix elements of the volume operator in canonical quantum gravity *J. Math. Phys.* **39** 3347–3371
- [14] Thiemann T 1998 A length operator for canonical quantum gravity *Jour. Math. Phys.* **39** 3372–3392
- [15] Major S A 1999 Operators for quantized directions *Class. Quant. Grav.* **16** 3859–3877
- [16] Thiemann T 1996 Anomaly-free formulation of non-perturbative, four-dimensional Lorentzian quantum gravity *Phys. Lett.* **B380** 257–264
- [17] Thiemann T 1998 Quantum spin dynamics (QSD) *Class. Quant. Grav.* **15** 839–873
- [18] Thiemann T. 1998 QSD III: Quantum constraint algebra and physical scalar product in quantum general relativity *Class. Quant. Grav.* **15** 1207–1247
- [19] Thiemann T 2001 *Modern Canonical Quantum General Relativity*. Cambridge: Cambridge University Press, in press, [gr-qc/0110034]
Ashtekar A, Lewandowski J 2004 Background independent quantum gravity: A status

- report *Class. Quant. Grav.* **21**, R53 [gr-qc/0404018]
 Smolin L 2004 An invitation to loop quantum gravity [hep-th/0408048]
 Rovelli C. (2004): *Quantum Gravity*, Cambridge: Cambridge University Press, 2004
- [20] Thiemann T 1998 QSD V: Quantum gravity as the natural regulator of matter quantum field theories *Class. Quant. Grav.* **15** 1281–1314
 Thiemann T 1998 Kinematical Hilbert Spaces for Fermionic and Higgs Quantum *Class. Quant. Grav.* **15** 1487-1512 [gr-qc/9705021]
- [21] Ashtekar A, Lewandowski J, Sahlmann H 2003 Polymer and Fock representations for a Scalar field *Class. Quant. Grav.* **20** L11-L21 [gr-qc/0211012]
- [22] Kamiński W, Lewandowski J, Bobieński B 2006 Background independent quantizations: the scalar field I *Class. Quant. Grav.* **23** 2761-2770
- [23] Lewandowski J, Okołów A, Sahlmann H and Thiemann T 2005 Uniqueness of diffeomorphism invariant states on holonomy-flux algebras *Preprint* gr-qc/0504147
- [24] Schmudgen K 1990 *Unbounded Operator Algebras and Representation Theory* (Basel-Boston-Berlin: Birkhauser Verlag)
- [25] Simon B 1999 *The Classical Moment Problem as a Self-Adjoint Finite Difference Operator* math-ph/9906008
- [26] Łojasiewicz S 1964 Triangulation of semi-analytic sets *Ann. Scuola. Norm. Sup. Pisa* 18 449-74
 Bierstone E and Milman P D 1988 Semianalytic and subanalytic sets *Publ Maths. IHES* 67 5-42
- [27] Fleischhack C 2004 Representations of the Weyl algebra in quantum geometry [math-ph/0407006]